

เอกสารอ้างอิง (ถ้าเกี่ยวข้องกับความดันโลหิตสูง)

1. Takada K, Sugiyama K, Yamamoto I. Protein hydrolyzates as antihypertensives. Patent: Jpn Kokai Tokkyo Koho JP 62,169,732 [87,169,732], 1987:10pp.
2. Matsudomi N, Kato A, Kobayashi K. Renin and angiotensin-converting enzyme inhibitors in food protein hydrolyzates. *Daizu Tanpakushitsu Eiyo Kenkyukai Kaishi* 1991;12(1):74-9.
3. Shimomura T, Yamada M, Asakura M, Kawamura K. Peptides as angiotensin-converting enzyme inhibitors and their manufacture with bromelain. Patent: Jpn Kokai Tokkyo Koho JP 04,299,991 [92,299,991], 1992:7pp.
4. Wu J, Ding X. Development of anti-hypertension peptide from soybean. I. Proteolytic enzyme from soybean as inhibitor of angiotensin converting enzyme I. *Zhongguo Youzhi* 1998;23(2):49-51.
5. Wu J, Ding X. Preparation of antihypertensive peptide from soybean: optimization of proteolytic conditions of enzyme E. *Zhongguo Youzhi* 1998;23(3):6-8.
6. Wu J, Ding X. Development of antihypertensive peptide from soybean: study on hydrolysis parameters of enzyme E. *Zhongguo Youzhi* 1998;23(4):12-3, 16.
7. Chen J-R, Okada T, Muramoto K, Suetsuna K, Yang S-C. Identification of angiotensin I-converting enzyme inhibitory peptides derived from the peptic digest of soybean protein. *Journal of Food Biochemistry* 2003;26(6):543-54.
8. Fan J, Saito M, Tatsumi E, Li L. Preparation of angiotensin I-converting enzyme inhibiting peptides from soybean protein by enzymatic hydrolysis. *Food Science Technology Research* 2003;9(3):254-6.
9. Matsui T. Production of hypotensive peptide, SVY, from 7S globulin of soybean and its physiological functions. *Daizu Tanpakushitsu Kenkyu* 2003;6:73-7.
10. Tanaka M, Kudo S, Watanabe T, Oikawa K, Uchita T, Kanazawa T. Angiotensin-converting enzyme inhibitors and their manufacture from soybean. Patent: Jpn Kokai Tokkyo Koho JP 2002 53,595, 2002:8pp.
11. Suetsuna K, Watanabe K. Isolation of new hexapeptides from soybean as inhibitor of angiotensin converting enzyme. Patent: Jpn Kokai Tokkyo Koho JP 2005139158, 2005:10pp.
12. Okamoto A, Hanagata H, Matsumoto E, Kawamura Y, Yanagida F. Antihypertensive substances in viscous material of fermented soybean (natto). *Food Hydrocolloids [Proc Int Conf Ind Exhib]* 1993:497-502.

13. Okamoto A, Hanagata H, Kawamura Y, Yanagida F. Anti-hypertensive substances in fermented soybean natto. *Plant Foods Hum Nutr* (Dordrecht, Neth) 1995;47(1):39-47.
14. Kuba M, Tanaka K, Tawata S, Takeda Y, Yasuda M. Angiotensin I-converting enzyme inhibitory peptides isolated from tofuyo fermented soybean food. *Biosci Biotechnol Biochem* 2003;67(6):1278-83.
15. Kataoka S. Functional effects of Japanese style fermented soy sauce (shoyu) and its components. *J Biosci Bioeng* 2005;100(3):227-34.
16. Zae-ik S, Rina Y, Soo-Ah P, et al. His-His-Leu, an angiotensin I converting enzyme inhibitory peptide derived from Korean soybean paste, exerts antihypertensive activity *in vivo*. *J Agric Food Chem* 2001;49:3004-9.
17. Yonekura M, Tanaka A. Isolation and application of physiologically active peptides from soybean whey and okara proteins. *Daizu Tanpakushitsu Kenkyu* 2003;6:88-93.
18. Tsurushiin S, Yoshida K, Nagano K, Fukuba H. Effect of heating on the ACE inhibitory activity of a crude soybean extract. *Nippon Kasei Gakkaishi* 1997;48(5):443-7.
19. Kasai N, Ikehara H. Stepwise extraction of proteins and carbohydrates from soybean seed. *J Agric Food Chem* 2005;53:4245-52.
20. Yu R, Park S-A, Chung D-K, Nam H-S, Shin Z-I. Effect of soybean hydrolyzate on hypertension in spontaneously hypertensive rats. *Han'guk Sikp'um Yongyang Kwahak Hoechi* 1996;25(6):1031-6.
21. Chen J-R, Yang S-C, Suetsuna K, Chao JC-J. Soybean protein-derived hydrolysate affects blood pressure in spontaneously hypertensive rats. *Journal of Food Biochemistry* 2004;28(1):61-73.
22. Yang H-Y, Yang S-C, Chen J-R, Tzeng Y-H, Han B-C. Soyabean protein hydrolyzate prevents the development of hypertension in spontaneously hypertensive rats. *Br J Nutr* 2004;92:507-12.
23. Wu J, Ding X. Hypotensive and physiological effect of angiotensin converting enzyme inhibitory peptides derived from soy protein on spontaneously hypertensive rats. *J Agric Food Chem* 2001;49(1):501-6.

24. Aoki H, Furuya Y, Endo Y, Fujimoto K. Effect of γ -aminobutyric acid-enriched tempeh-like fermented soybean (GABA-tempeh) on the blood pressure of spontaneously hypertensive rats. *Biosci Biotechnol Biochem* 2003;67(8):1806-8.
25. Shizuka F, Kido Y, Nakazawa T, Kitajima H, Aizawa C, Kayamura H, Ichijo N. Antihypertensive effect of γ -aminobutyric acid enriched soy products in spontaneously hypertensive rats. *Biofactors* 2004;22(1-4):165-7.
26. Martin DS, Breitkopf NP, Eyster KM, Williams JL. Dietary soy exerts an antihypertensive effect in spontaneously hypertensive female rats. *Am J Physiol* 2001;281(2, Pt 2):R553-60.
27. Martin DS, Williams JL, Breitkopf NP, Eyster KM. Pressor responsiveness to angiotensin in soy-fed spontaneously hypertensive rats. *Can J Physiol Pharmacol* 2002;80:1180-6.
28. Park E, Shin JI, Park OJ, Kang MH. Soy isoflavone supplementation alleviates oxidative stress and improves systolic blood pressure in male spontaneously hypertensive rats. *J Nutr Sci Vitaminol (Tokyo)* 2005;51(4):254-9.
29. Washburn S, Burke GL, Moargant T, Anthony M. Effect of soy protein supplementation on serum lipoproteins, blood pressure, and menopausal symptoms in perimenopausal women. *J Amer Menopause Soc* 1999;6(1):7-13.
30. Nagata C, Shimizu H, Takami R, Hayashi M, Takeda N, Yasuda K. Association of blood pressure with intake of soy products and other food groups in Japanese men and women. *Prev Med* 2003;36(6):692-7.
31. He J, Gu D, Wu X, et al. Effect of soybean protein on blood pressure: a randomized, controlled trial. *Ann Intern Med* 2005;143(1):1-9.
32. Rivas M, Garay RP, Escanero JF, Cia Jr P, Cia P, Alda JO. Soy milk lowers blood pressure in men and women with mild to moderate essential hypertension. *J Nutr* 2002;132:1900-2.
33. Javed A, Ali S, Shah WH, Salam A. Effect of dietary supplementation of fatty acids on various blood parameters of male essential hypertensive patients. *Journal of the Chemical Society of Pakistan* 2003;25(2):172-5.
34. Fujimaki A. Antihypertensive agents. Patent: Eur Pat Appl EP 78,028, 1983:30pp.
35. Nakamura T, Ota N. Proteins and fats as antihypertensives. Patent: Jpn Kokai Tokkyo Koho JP 2002 363,095, 2002:6pp.

36. Mizutani T, Ogoshi E, Miura K, Fujimoto Y. Soybean fermentation products for prevention of diabetes complication. Patent: Jpn Kokai Tokkyo Koho JP 2003 300,894, 2003.
37. Kasai N, Imai T, Nakai K. Preparation of protein-high soybean extract. Patent: Jpn Kokai Tokkyo Koho JP 2005095156, 2005:13pp.
38. Moriyama T, Meabuchi M, Maruyama N. Soybean protein β -conglycinin-containing food materials, food, and pharmaceuticals for treatment of life-style diseases and inhibition of fatty acid synthase activity. Patent: Jpn Kokai Tokkyo Koho JP 2005080533, 2005:11pp.
39. Matoba N, Doyama N, Yamada Y, Maruyama N, Utsumi S, Yoshikawa M. Design and production of genetically modified soybean protein with antihypertensive activity by incorporating potent analog of ovokinin(2-7). FEBS Lett 2001;497(1):50-4.
40. Matoba N, Yamada Y, Yoshikawa M. Design of a genetically modified soybean protein preventing hypertension based on an anti-hypertensive peptide derived from ovalbumin. Curr Med Chem Cardiovasc Hematol Agents 2003;1(2):197-202.
41. Matoba N, Doyama N, Yamada Y, Maruyama N, Utsumi S, Yoshikawa M. Design and production of genetically modified soybean protein with anti-hypertensive activity by incorporating potent analogue of ovokinin(2-7).
42. Onishi K, Matoba N, Yamada Y, Doyama N, Maruyama N, Utsumi S, Yoshikawa M. Optimal designing of β -conglycinin to genetically incorporate RPLKPW, a potent anti-hypertensive peptide. Peptides 2004;25(1):37-43.

เอกสารอ้างอิง (หมอกบัว)

1. Samitinand T. Thai Plant Name. Bangkok:The Forest Herbarium, Royal Department, 2001. 810+III pp.
2. Larsen K. DROSERACEAE. In: Smitinand T, Larsen K, eds. Flora of Thailand, Vol. 5 Part 1. Bangkok:The Chutima Press, 1987: 67-9.
3. วงศ์สฤติย์ ฉั่วกุล สมภพ ประธานธรรารักษ์ นพมาศ สุนทรเจริญนนท์ และคณะ. สมุนไพรพื้นบ้านอีสาน. ใน: วงศ์สฤติย์ ฉั่วกุล, บรรณาธิการวิชาการ. สารานุกรมสมุนไพร เล่ม 4 กกายอีสาน. กรุงเทพมหานคร:อมรินทร์พริ้นติ้งแอนด์พับลิชชิ่ง จำกัด (มหาชน), 2543:272 หน้า.
4. Duke JA, Ayensu ES. Medicinal plants of China. Michigan:Alconac, Reference Publications, Inc., 1985;1(4):52-361
5. Culham A, Gornall RJ. The taxonomic significance of naphthoquinones in the Droseraceae. Biochem Syst Ecol 1993;22(5):507-15.
6. Zeng X, Han J. Plumbagin-containing acaricide and its preparation process. Faming Zhuanli Shenqing Gongkai Shuomingshu 2004:14pp.
7. Samy RP, Ignacimuthu S, Sen A. Screening of 34 Indian medicinal plants for antibacterial properties. J Ethnopharmacol 1998;62(2):173-82.
8. Zenk MH, Furbringer M, Steglich W. Occurrence and distribution of 7-methyljuglone and plumbagin in the Droseraceae. Phytochemistry 1969;8:2199-200.
9. Kapur SK. Medico-botanic survey of medicinal and aromatic plants of mawpliang (shillong). Indian Drugs 1983;21(1):1-5.
10. Joshi DN, Sah BCL, Suri RK. Some medicinal plants of Rudranath bugyal (distt. Chamoli), U.P. Bull Med Ethnobot Res 1982;3(1):27-42.
11. Manandhar NP. Ethnobotany of Jumla district, Nepal. Int J Crude Drug Res 1986;24(2):81-9.
12. Asano M, Hase J. Hydroxyquinones. X. pigment of *Drosera peltata*. Yakugaku Zasshi 1943;63:410-1.
13. Liang W-J, Chang W-S. Extraction and assay of plumbagin from *Drosera peltata*. Yaoxue Tongbao 1980;15(8):3-4.
14. Ramachandran Nair AG, Shanmugasundaram P, Madhusudhanan KP. Naphthaquinones and flavonols from leaves of *Drosera peltata*. Fitoterapia 1990;61(1):85-6.

15. Hu XB, Yang PQ, Liu WJ. A new indanone isolated from *Drosera peltata* var. *lunata*. Yunnan Zhiwu Yanjiu 1991;13(3):334-40.
16. Liang WJ, Chang WS. Extraction and assay of plumbagin from *Drosera peltata*. Yao Hsueh T'ung Pao 1980;15(8):3-4.
17. Wang QA, Su JY, Zeng LM. Study on chemical constituents of *Drosera peltata* Smith var. *lunata* collected in Tibet. Zhongguo Zhongyao Zazhi 1998;23(11):683-4.
18. Didry N, Dubreuil L, Trotin F, Pinkas M. Antimicrobial activity of aerial parts of *Drosera peltata* Smith on oral bacteria. J Ethnopharmacol 1998;60(1):91-6.
19. Krenn L, Blaeser U, Hausknot-Chenicek N. Determination of naphthoquinones in *Droserae herba* by reversed-phased high performance liquid chromatography. J Liq Chromatogr 1998;21(20):3149-60.
20. Leclercq J, Angenot L. *Drosera peltata* and the standardization of *Drosera* tincture. Journal de Pharmacie de Belgique 1984;39(5):269-74.
21. Amagase S, Mori M, Nakayama S. Digestive enzymes in insectivorous plants. IV. Enzymeic digestion of insects by *Nepenthes* secretion and *Drosera peltata* extract. Proteolytic and chitinolytic activities. J Biochemistry 1972;72(3):765-7.
22. Amagase S. Digestive enzymes in insectivorous plants. III. Acid proteases in the genus *Nepenthes* and *Drosera peltata*. J Biochemistry 1972;72(1):73-81.
23. Van Der Vijver LM, Loetter AP. Constituents in the roots of *Plumbago auriculata* and *Plumbago zeylanica* responsible for antibacterial activity. Planta Med 1971;20:8.
24. Van Der Vijver LM. Distribution of plumbagin in the Plumbaginaceae. Phytochemistry 1972;11:3247.
25. Chowdhury AKA, Chakder SK, Khan AKA. Isolation and characterization of chemical constituents of *Plumbago zeylanica* root. J Bangladesh Acad Sci 1981;5(1):71-4.
26. Ko K. On the pharmacological action of plumbagin. Jap J Med Sci 4 Pharmacol 1933;6:259.
27. Durga R, Sridhar P, Polasa H. Antimutagenic activity of plumbagin in Ames *Salmonella typhimurium* test. Indian J Med Res 1992;96(2):143-5.
28. Anon. Ayurvedic drug to fight cancer. Probe 1985;24(4):234.

29. Kavimani S, Ilango R, Madheswaran M, Jayakar B, Gupta M, Majumdar UK. Antitumour activity of plumbagin against Dalton's ascitic lymphoma. *Indian J Pharm Sci* 1996;58(5):194-6.
30. Oommen E, Shenoy BD, Udupa N, Kamath R, Devi PU. Antitumor efficacy of cyclodextrin-complexed and niosome-encapsulated plumbagin in mice bearing melanoma B16F1. *Pharm Pharmacol Commun* 1999;5(4):281-5.
31. Singh UV, Bisht KS, Rao S, Uma DP, Udupa N. Plumbagin-loaded plga microspheres with reduced toxicity and enhanced antitumor efficacy in mice. *PHARM SCI* 1996;2(9):407-9.
32. Raja Naresh RA, Udupa N, Uma Devi P. Niosomal plumbagin with reduced toxicity and improved anticancer activity in BALB/C mice. *J Pharm Pharmacol* 1996;48(11):1128-32.
33. Krishnaswamy M, Purushothaman KK. Plumbagin: a study of its anticancer. Antibacterial and antifungal properties. *Indian J Exp Biol* 1980;18:876-7.
34. Chandrasekaran B, Nagarajan B. Growth inhibitory effect of echitamine and plumbagin on fibrosarcoma in rats. *Arogya* 1983;9(1):60-3.
35. Gujar GT. Plumbagin, a naturally occurring naphthoquinone. Its pharmacological and pesticidal activity. *Fitoterapia* 1990;61(5):387-94.
36. Kuo YH, Chang CI, Li SY, Chou CJ, Chen CF, Kuo YH, Lee KH. Cytotoxic constituents from the stems of *Diospyros maritima*. *Planta Med* 1997;63(4):363-5.
37. Moreira LC, Ferreira De Santana C, Nunes Da Silva I. In vitro study of the antineoplastic action of substances of microbial and plant origin against Ehrlich and Sarcoma 180 carcinomas. *Rev Inst Antibiot Univ Fed Pernambuco Recife* 1973;13:77-84.
38. Noda Y, Kawazoe Y, Hakura A. Cytotoxicity of naphthoquinones toward cultured resting murine leukemia L1210 cells in the presence of glutathione, diethyl maleate, or iodoacetamide. *Biol Pharm Bull* 1997;20(12):1250-6.
39. Fujii N, Yamashita Y, Arima Y, Nagashima M, Nakano H. Induction of topoisomerase II-mediated DNA cleavage by the plant naphthoquinones plumbagin and shikonin. *Antimicrob Agents Chemother* 1992;36(12):2589-94.
40. Bhatta BB, Lal S. The pharmacological action of *Plumbago zeylanica* and its active principle (plumbagin). *Indian J Med Res* 1933;20:777-88.

41. Debray M, Araud O, Paris RR. A native Madagascan Ebenaceae, *Diospyros gracilipes*. *Plant Med Phytother* 1973;7:77.
42. Tummin Katti MC, Patwardhan VN. Chemical examination of the root bark of *Plumbago rosea*. *J Indian Inst Sci Ser A* 1932;15:9-15 ()
43. Ragazzi,E: De Biasi M, Pandolfo L, Chinellato A, Caparrotta L. In vitro effects of naphthoquinones isolated from *Drosera* species. *Pharmacol Res* 1993;27(1):87-8.
44. Premakumari P, Rathinam K, Santhakumari G. Antifertility activity of plumbagin. *Indian J Med Res* 1977;65:829-38.
45. Bhargava SK, Dixit VP. Antifertility effects of embelin and plumbagin in female rats. *Plant Med Phytother* 1985;19(1):29-34.
46. Santhakumari C, Suganthan D, Pillai NR, Seshadri C, Nair RB. Effect of plumbagin on spermatogenesis and accessory sex organs in adult rats. *Indian J Pharmacol* 1981;13(1):66.
47. Bhargava SK. Effects of plumbagin on reproductive function of male dog. *Indian J Exp Biol* 1984;22:153-6.
48. Vichkanova SA, Adgina VV, Goryunova LV, Rubinchik MA. Chemotherapeutic properties of plumbagin. *Fitontsidy Mater Soveshch* 6th 1972:183.
49. Tikkanen L, Matsushima T, Natori S, Yoshihira K. Mutagenicity of natural naphthoquinones and benzoquinones in the *Salmonella*/microsome test. *Mutat Res* 1983;124(1):25-34.