

- (1) Fu C, Shi H, Li, Q. A review on pharmacological activities and utilization technologies of pumpkin. *Plant Foods Hum Nutr* 2006; 61: 73-80.
- (2) Ribaya-Mercado JD, Maramag CC, Tengco LW, Dolnikowski GG, Blumberg JB, Solon FS. Carotene-rich plant foods ingested with minimal dietary fat enhance the total-body vitamin A pool size in Filipino schoolchildren as assessed by stable-isotope-dilution methodology. *Am J Clin Nutr* 2007; 85: 1041-1049.
- (3) Huang X-E, Hirose K, Wakai K, Matsuo K, Ito H, Xiang J, Takezaki T, Tajima K. Comparison of lifestyle risk factors by family history for gastric, breast, lung and colorectal cancer. *Asian Pac J Cancer Prev* 2004; 5(4): 419-427.
- (4) Colditz GA, Branch LG, Lipnick RJ, Willett WC, Rosner B, Posner BM, Hennekens CH. Increased green and yellow vegetable intake and lowered cancer deaths in an elderly population. *Am J Clin Nutr* 1985; 41: 32-36.
- (5) Goodman GE, Thornquist MD, Balmes J, Cullen MR, Meyskens FL Jr, Omenn GS, Valanis B, Williams JH Jr. The beta-carotene and retinol efficacy trial: incidence of lung cancer and cardiovascular disease mortality during 6-year follow-up after stopping beta-carotene and retinol supplements. *J Natl Cancer Inst* 2004; 96 (23): 1743-1750.
- (6) Lee DH, Iwanski GB, Thoennissen NH. Cucurbitacin: Ancient compound shedding new light on cancer treatment. *Scientific World Journal* 2010; 10: 413-418.
- (7) Xiaomin H, Edward JM, Jieming X, Mingdong H, Minghuang C, Liqing C. Atomic resolution structure of cucurmosin, a novel type 1 ribosome-inactivating protein from the sarcocarp of *Cucurbita moschata*. *J Struct Biol* 2008; 164: 81-87.
- (8) Villaseñor IM, Domingo AP. Anticarcinogenicity potential of spinasterol isolated from squash flowers. *Tertog Carcinog Mutagen* 2000; 20 (3): 99-105.
- (9) Paiva SA, Russell RM. Beta-carotene and other carotenoids as antioxidants. *J Am Coll Nutr* 1999; 18(5): 426-433.
- (10) Xanthopoulou MN, Nomikos T, Fragopoulou E, Antonopoulou S. Antioxidant and lipoxygenase inhibitory activities of pumpkin seed extracts. *Food Res Int* 2009; 42: 641-646.
- (11) Yang X, Zhao Y, Lv Y. Chemical composition and antioxidant activity of an acidic polysaccharide extracted from *Cucurbita moschata* Duchesne ex Poiret. *J Agric Food Chem* 2007; 55 (12): 4684-4690.
- (12) Makni M, Fetoui H, Gargouri NK, Garoui EM, Jaber H, Makni J, et al. Hypolipidemic and hepatoprotective effects of flax and pumpkin seed mixture rich in ω -3 and ω -6 fatty acids in hypercholesterolemic rats. *Food Chem. Toxicol.* 2008; 46: 3714-3720.
- (13) Li QH, Fu CL, Rui YK, Hu GH, Cai TY. Effects of protein-bound polysaccharide isolated from pumpkin on insulin in diabetic rats. *Plant Foods Hum Nutr* 2005; 60: 13-16.

- (14) Li QH, Tian Z, Cai TY. Study on the hypoglycemic action of pumpkin extract in diabetic rat. *Acta Nutrimenta Sin* 2001; 25 (1): 34-36.
- (15) Friederic M, Theurer C, Schiebel-Schlosser G. Prosta Fink Forte capsules in the treatment of benign prostatic hyperplasia. Multicentric surveillance study in 2245 patients. *Forsch Komplementarmed Klass Naturheilkd* 2000; 7 (4): 200-204.
- (16) Dreikorn K. The role of phytotherapy in treating lower urinary tract symptoms and benign prostatic hyperplasia. *World J Urol* 2002; 19 (6): 426-435.
- (17) Hammer KA, Carson CF, Riley TV. Antimicrobial activity of essential oils and other plant extracts. *J Appl Microbiol* 1999; 86 (6): 985-990.
- (18) Villaseñor IM, Bartolome ALO, Daclan LC, Manatad AJG. Microbiological and pharmacological studies on extracts of *Cucurbita maxima*. *Phytother Res* 1995; 9: 376-378.
- (19) Jun H-II, Lee CH, Song GS, Kim YS. Characterization of the pectic polysaccharides from pumpkin peel. *LWT-Food Science and Technology* 2006; 39 (5): 554-561.
- (20) Park S-C, Kim J-Y, Lee J-K, Hwang I, Cheong H, Nah J-W, Hahm K-S, Park Y. Antifungal mechanism of a novel antifungal protein from pumpkin rinds against various fungal pathogens. *J Agric Food Chem* 2009; 57 (19): 9299-9304.
- (21) Cheong NE, Choi YO, Kim WY, Bae IS, Cho MJ, Hwang I, Kim JW, Lee SY. Purification and characterization of an antifungal PR-5 protein from pumpkin leaves. *Mol Cells* 1997; 7 (2): 214-219.
- (22) Wang HX, Ng TB. Isolation of cucurmoschin, a novel antifungal peptide abundant in arginine, glutamate and glycine residues from black pumpkin seeds. *Peptides* 2003; 24: 969-972.
- (23) Vassiliou AG, Neumann GM, Condron R, Polya GM. Purification and mass spectrometry-assisted sequencing of basic antifungal proteins from seeds of pumpkin (*Cucurbita maxima*). *Plant Sci* 1998; 134: 141-162.
- (24) Elisha EE, Twajj AA, Ali NM, Tarish JH, Al-Omari MM, Karim S. The anthelmintic activity of some Iraqi plants of the Cucurbitaceae. *Int J Crude Drug Res* 1987; 25: 153-157.
- (25) Diaz OD, Lloja LL, Carbajal ZV. Preclinical studies of *Cucurbita maxima* (pumpkin seeds) a traditional intestinal antiparasitic in rural urban areas. *Res Gastroentero Peru* 2004; 24 (4): 323-327.
- (26) นิตยสารหมอชาวบ้าน ฟักทอง: ยาขับพยาธิ 2529; 6 (69): 43-44.
- (27) Suphiphat V, Morjaroen N, Pukboonme I, Ngumboonsri P, Lowhnoo T, Dhanamitta S. The effect of pumpkin seeds snack on inhibitors and promoters of urolithiasis in Thai adolescents. *J Med Assoc Thai* 1993; 76 (9): 487-493.
- (28) Yun HS, Chang IM. Plants with liver protective activities. *Kor J Pharmacog* 1977; 8 (3): 125-129.

เอกสารอ้างอิง Argan oil ๒๗ (๔)

1. Maurin R. Argan oil, *Argania spinosa* (L.) Skeels (Sapotaceae). *Revue Francaise des Coeps Gras* 1992;39(5-6):139-46.
2. Khallouki F, Spiegelhalter B, Bartsch H, Owen RW. Secondary metabolites of the argan tree (Morocco) may have disease prevention properties. *African J Biotech* 2005;4(5):381-8.
3. Charrouf Z, Guillaume D. Ethnoeconomical, ethnomedical, and phytochemical study of *Argania spinosa* (L.) Skeels. *J Ethnopharmacology* 1999;67:7-14.
4. Charrouf Z, Dube S. Helping Moroccan women preserve the argan tree at the gateway to the Sahara. *IDRC Bulletin* 2000. (<http://www.idrc.org/en/ev-5267-201-2-DO-TOPIC.html>)
5. Nerd A, Eteshola E, Borowy N, Mizrahi Y. Growth and oil production of argan in the Negev desert of Israel. *Industrial Crops and Products* 1994;2(2):89-95.
6. Belardi-Benmahdi M M, Khaldi D, Baghdad C, Gouzi H, Bendimerad N, Hammonti B. Physicochemical and nutritional study of argan oil (*Argania spinosa* L.) in south-western Algeria. *Pigment & Resin Technology* 2009;38(2):96-9.
7. Huyghebaert A, Hendrickx H. Chemical, physical, and technological aspects of argan oil. *Oleagineux* 1974;29(1):29-31.
8. Charrouf Z, Guillaume D. Argan oil: occurrence, composition and impact on human health. *Eur J Lipid Sci Technol* 2008;110(7):632-36.
9. Sasserath EA. "Morocco Olive oil". *Zeitschrift fuer Untersuchung der Nahrungs-und Genussmittel Sowie de Gebrauchsgegenstaende* 1911;20:749-50.
10. Charrouf Z. "Valorisation d'Argania spinosa (L.) Sapotaceae: Etude de la composition Clinique et de l'activite biologique du tourteau et de l'extrait lipidique de la pulpe". These Sciences, Univ Mohammed V, Rabat, 1991.
11. Charrouf Z, Guillaume D. Argan oil, functional food and the sustainable development of the argan forest. *Natural Product communications* 2008b;3(2):283-8.
12. Girardet A. Some constants of argan oil. *Pharmaceutica Acta Helvetiae* 1927;2:1-5.
13. Belcadi-Haloui R, Zekhni A, Hatimi A. Comparative study on fatty acid and tocopherol composition in argan oils extracted from fruits of different forms. *Acta Botanica Gallica* 2008;155(2):301-5.
14. Domokos J, Kiss B, Budapest H. A rarely used dietetic and cosmetic oil: argan oil. *Olaj, Szappan, Kozmetika* 2007;56(3):100-3.

15. Yaghmur A, Aserin A, Mizrahi Y, Nerd A, Garti N. Evaluation of argan oil for deep-frying. *Lebensmittel-Wissenschaft und-Technologie* 2001;34(3):124-30.
16. Seidermann J. Falsification of seed oil of *Argania spinosa*. *Deutsche Lebensmittel-Rundschau* 1998;94(1):26-7.
17. Alaoui BF, Zerriouh A, Belkbir L. Conservation study of the Argan oil by thermogravimetry. *Chemistry* 2001;13(1):144-50.
18. Hammadi C, Cillard J, Cillard P. Autoxidation of argan oil, *Argania spinosa* L. from Morocco. *Sciences des Aliments* 1994;14(1):117-24.
19. Khallouki F, Mannina L, Viel S, Owen RW. Thermal stability and long –chain fatty acid positional distribution on glycerol of argan oil. *Food Chemistry* 2008;110(1):57-61.
20. Khallouki F, Younos C, Soulimani R, Oster T, Charrouf Z, Spiegelhalder B, Bartsch H, Owen RW. Consumption of argan oil (Morocco) with its unique profile of fatty acids, tocopherols, squalene, sterols and phenolic compounds should confer valuable cancer chemopreventive effects. *European Journal of Cancer Prevention* 2003;12(1):67-75.
21. Cayuela JA, Rada M, Perez-Camino MC, Benaissa M, Abdelaziz E, Guida A. *European Journal of Lipid Science and Technology* 2008;110(12):1159-66.
22. Farines M, Soulier J, Charrouf M, Cave A. Etude de l'huile des graines d'*Argania spinosa* (L.); Sapotaceae. II Sterols, alcools, triterpeniques et methylsterols de l'huile d'argan. *Revue Francaise des Caps Gras* 1984;31:443-8.
23. Rojas LB, Quideau S, Pardon P, Charrouf Z. Colorimetric evaluation of phenolic content and GC-MS characterization of phenolic composition of alimentary and cosmetic argan oil and press cake. *Journal of Agricultural and Food Chemistry* 2005;53(23):9122-7.
24. Zahar M, Reineccius G, Schirle-Keller J-P. Identification of aroma compounds in food grade Argan oil. Abstracts of papers, 234th ACS National Meeting, Boston, MA, United States, August 19-23, 2007 (2007), AGFD-072.
25. Marfil R, Caberra-Vique C, Gimenez R, Bouzas PR, Martinez O, Sanchez JA: Metal content and physicochemical parameters used as quality criteria in virgin argan oil: influence of the extraction method. *J Agri Food Chem* 2008;56(16):7279-84.
26. Cherki M, Berrougui H, Drissi A, Adlouni A, Khalil A. Argan oil: which benefits on cardiovascular diseases? *Can Pharmacol Res* 2006;54(1):1-5.

27. Mekhfi H, Gadi D, Bnouham M, Ziyyat A, Legssyer A, Aziz M. Effect of argan oil on platelet aggregation and bleeding time: a beneficial nutritional property. *J Complementary and Integrative Medicine* 2008;5(1): (no pp given).
28. Berrougui H, Cloutier M, Isabelle M, Khalil A. Phenolic-extract from argan oil (*Argania spinosa* L.) inhibits human low-density lipoprotein (LDL) oxidation and enhances cholesterol efflux from human THP-1 macrophages. *Atherosclerosis* 2006;184(2):389-96.
29. Berrougui H, Alvarez de Sotomayor M, Perez-Guerrero C, Ettaib A, Hmamouchi M, Marhuenda E, Herrera MD. Argan (*Argania spinosa*) oil lowers blood pressure and improves endothelial dysfunction in spontaneously hypertensive rats. *Bri J Nutr* 2004;92(6):921-9.
30. Berrougui H, Ettaib A, Herrera Gonzalez MD, Alvarez de Solomayor M, Bennani-Kabchi N, Hmamouchi M. Hypolipidemic and hypocholesterolemic effect of argan oil (*Argania spinosa* L.) in Meriones Shawi rats. *J Ethnopharmacol* 2003;89(1):15-8.
31. Benajiba N, Morel S, De Leiris J, Boucher F, Charrouf Z, Mokhtar N, Aguenau H. The effect of argan oil on heart function during ischemia and reperfusion. *Therapie* 2002;57(3):246-52.
32. Benajiba N, De Leiris J, Lyan B, Derouiche A, Mokhtar N, Aguenau H. Effect of argan oil on aortic contractility:susceptibility to oxidative stress. *Oleagineux, Corps Gras, Lipises* 2006;13(1):76-80.
33. Drissi A, Girona J, Cherki M, Godas G, Derouiche A, El Messal M, Saile R, Kettani A, Sola R, Masana L, Adlouni A. Evidence of hypolipemiant and antioxidant properties of argan oil derived from the argan tree (*Argania spinosa*). *Clin Nutr* 2004;23(5):1159-66.
34. Derouiche A, Cherki M, Drissi A, Bamou Y, El Messal M, Idrissi-Oudghiri A, Lecerf JM, Adlouni A. Nutritional intervention study with argan oil in man:effect on lipids and lipoproteins. *Annals of Nutrition & Metabolism* 2005;49(3):196-201.
35. Cherki M, Derouiche A, Drissi A, El Messal M, Bamou Y, Idrissi-Quadghiri A, Khali A, Adlouni A. Consumption of argan oil may have an antiatherogenic effect by improving peraoxonase activitites and antioxidant status: Intervention study in healthy men. *Nutrition, Metabolism, and cardiovascular diseases* 2005;15(5):352-60.
36. Bennani H, Drissi A, Giton F, Kheuang L, Fiet J, Adlouni A.. Antiproliferative effect of polyphenols and sterols of virgin argan oil on human prostate cancer cell lines. *Cancer Detection and Prevention* 2007;31(1):64-9.

37. Drissi A, Bennani H, Giton F, Charrouf Z, Fiet J, Adlouni A. Tocopherols AND SAPONINS DERIVED FROM *Argania spinosa* Exert, and antiproliferative effect on human prostate cancer. *Cancer Investigation* 2006;24(6):588-92.
38. Samane S, Noel J, Charrouf Z, Amarouch H, Haddad PS. Insulin-sensitizing and anti-proliferative effects of *Argania spinosa* seed extracts. *Evidence-based Complementary and Alternative Medicine* 2006;3(3):317-27.
39. Samane S, Christon R, Dombrowski L, Turcotte S, Charrouf Z, Lavigne C, Levy E, Bachelard H, Amarouch H, Marette A, Haddad PS. Fish oil and argan oil intake differently modulate insulin resistance and glucose intolerance in a rat model of dietary-induced obesity. *Metabolism, Clinical and Experimental* 2009;58(7):909-19.
40. Benzaria A, Meskini N, Dubois M, Croset M, Nemoz G, Lagarde M, Prigent AF. Effect of dietary argan oil on fatty acid composition, proliferation, and phospholipase D activity of rat thymocytes. *Nutrition* 2006;22(6):628-37.