

เอกสารอ้างอิง ชะเอมเทศ

1. British Herbal Medicine Association. British Herbal Pharmacopoeia. Bournemouth: Megaron Press, 1983.
2. World Health Organization. WHO Monographs on Selected Medicinal Plants. Vol. 1. Geneva: World Health Organization, 1999.
3. Government of India, Ministry of Health and Family Welfare. The Ayurvedic Pharmacopoeia of India. Part-1. 1st ed. Vol. 1. Delhi: Government of India, Ministry of Health and Family Welfare, Department of India Systems of Medicine & Homoeopathy, 2001.
4. Karnick CR. Pharmacopoeial Standards of Herbal Plants. Vol. 1. Delhi: Sri Satguru Publications, 1994.
5. China Pharmacopoeia Commission. Pharmacopoeia of The People's Republic of China. Vol. 1. Beijing: People's Republic of China, 2005.
6. Yeung HC. Handbook of Chinese Herbs (Chinese Materia Medica). California: Los Angeles County, 1996.
7. The Ministry of Health and Welfare. Pharmacopoeia of Japan. 14th ed. (English version). Tokyo: The Ministry of Health and Welfare, 2001.
8. Blumenthal M, Goldberg A, Brinckmann J. Herbal Medicine expanded Commission E Monographs. MA: Integrative Medicine Communications, 2000.
9. Ulbricht CE, Basch EM (eds). Natural Standard Herb & Supplement Reference Evidence-based Clinical Reviews. Missouri: Elsevier Mosby, 2005.
10. The Forest Herbarium, Royal Forest Department. Thai Plant Names Tem Smitinand. Revised ed. Bangkok: Prachachon Co., 2001.
11. Bisset NG. Herbal Drugs and Phytopharmaceuticals. London: CRC Press, 1994.
12. Nagai H, Yamamoto Y, Sato Y, Akao T, Tani T. Pharmaceutical evaluation of cultivate *Glycyrrhiza uralensis* roots in comparison of their antispasmodic activity and glycycomarin contents with those of licorice. Bio Pharm Bull 2006; 29(12): 2442-5.
13. Tang W, Eisenbrand G. Chinese Drugs of Plant Origin. Berlin: Springer-Verlag, 1992.
14. Bai H, Li W, Koike K, Dou D, Pei Y, Chen Y, Nikaido T. A novel biflavonoid from roots of *Glycyrrhiza uralensis* cultivated in China. Chem Pharm Bull 2003; 51(9): 1095-7.

15. Nomura T, Fukai T, Akiyama T. Chemistry of phenolic compounds of licorice (*Glycyrrhiza* species) and their estrogenic and cytotoxic activities. *Pure Appl Chem* 2002; 74(7): 1199-206.
16. Fukai T, Tantal L, Nomura T. Isoprenoid-substituted flavanoids from *Glycyrrhiza glabra*. *Phytochemistry* 1996; 43(2): 531-2.
17. Fukai T, Cai BS, Maruno K, Horikoshi T, Nomura T. Isoprenylated flavanoids from underground parts of *Glycyrrhiza glabra*. *Phytochemistry* 1996; 43(5): 1119-24.
18. Fukai T, Cai BS, Maruno K, Miyakawa Y, Konishi M, Nomura T. An isoprenylated flavanone from *Glycyrrhiza glabra* and rec-assay of licorice phenols. *Phytochemistry* 1998; 49(7): 2005-13.
19. Tsukamoto S, Aburatani M, Yoshida T, Yamashita Y, El-Beih AA, Ohta T. CYP3A inhibitors isolated from licorice. *Bio Pharm Bull* 2005; 28(10): 2000-2.
20. He J, Chen L, Heber D, Shi W, Lu QY. Antibacterial compounds from *Glycyrrhiza uralensis*. *J Nat Prod* 2006; 69(1): 121-4.
21. ชัยนนต์ พิเชียรสุนทร แม้นมาส ชาลิต วิเชียร จีรวงศ์. คำอธิบายตำราพระโอสถพระนารายณ์. กรุงเทพมหานคร: สำนักพิมพ์อมรินทร์, 2542.
22. Burnham TH (ed). *The Review of Natural Products*. 1rd ed. Missouri: Facts and Comparisons, 2001.
23. Mae T, Kida H, Nishiyama T, Tsukagawa M, Konishi E, Kuroda M, Mimaki Y, Sashida Y, Takahashi K, Kawada T, Nakagawa K, Kitahara M. A licorice ethanolic extract with peroxisome proliferators-activated receptor-gamma ligand-binding activity affects diabetes in KK-Ay mice, abdominal obesity in diet-induced obese C57BL mice and hypertension in spontaneously hypertensive rats. *J Nutr* 2003; 133(11): 3369-77.
24. Kim DC, Choi SY, Kim SH, Yun BS, Yoo ID, Reddy NR, Yoon HS, Kim KT. Isoliquiritigenin selectively inhibits H(2) histamine receptor signaling. *Mol Pharmacol* 2006; 70(2): 493-500.
25. Sato Y, He JX, Nagai H, Tani T, Akao T. Isoliquiritigenin, one of the antispasmodic principles of *Glycyrrhiza uralensis* roots, acts in the lower part of intestine. *Biol Pharm Bull* 2007; 30(1): 145-9.
26. Sato Y, Akao T, He JX, Nojima H, Kuraishi Y, Morota T, Asno T, Tani T. Glycycoumarin from *Glycyrrhizae Radix* acts as a potent antispasmodic through inhibition of phosphodiesterase. *J Ethnopharmacol* 2006; 105(3): 409-14.

27. Ahn J, Um M, Choi W, Kim S, Ha T. Protective effects of *Glycyrrhiza uralensis* Fisch. on the cognitive deficits caused by beta amyloid peptide 25-35 in young mice. *Biogerontology* 2006; 7(4): 239-47.
28. Takahashi T, Takasuka N, Iigo M, Nishino H, Tsuda H, Okuyama T. Isoliquiritigenin, a flavonoid from licorice, reduces prostaglandin E₂ and nitric oxide, cause apoptosis, and suppresses aberrant crypt foci development. *Cancer Sci* 2004; 95(5): 448-53.
29. Jo EH, Hong HD, Ahn NC, Jung JW, Yang SR; Park JS, Kim SH, Lee YS, Kang KS. Modulations of the Bcl-2/Bax family were involved in the chemopreventive effects of licorice root (*Glycyrrhiza uralensis* Fisch.) in MCF-7 human breast cancer cell. *J Agric Food Chem* 2004; 52(6): 1715-9.
30. Fukai T, Satoh K, Nomura T, Sakagami H. Antinephritis and radical scavenging activity of prenylflavonoids. *Fitoterapia* 2003; 74(7-8): 720-4.
31. Jo EH, Kim SH, Ra JC, Kim SR, Cho SD, Jung JW, Yang SR; Park JS, Hwang JW. Chemopreventive properties of the ethanol extract of Chinese licorice (*Glycyrrhiza uralensis*) root: induction of apoptosis and G1 cell cycle arrest in MCF-7 human breast cancer cells. *Cancer Lett* 2005; 230(2): 239-47.