

เอกสารอ้างอิง นำ้สูกยอ (โนนิ) ต้านมะเร็งได้จริง?

1. พร้อมจิต ศรลัมพ์ รุ่งระวี เต็มศิริกนกุล วงศ์สติตย์ จั่วกุล และคณะ. สมุนไพรสวนสิรีรุกขชาติ. กรุงเทพฯ: บริษัทอมรินทร์พรินติ้งกรุ๊ฟ จำกัด, 2535:257 หน้า.
2. นันทวน บุณยะประภัคร อรุณช โชคชัยเจริญพร (บรรณาธิการ). สมุนไพรไม้พื้นบ้าน (4). กรุงเทพฯ: บริษัทประชาชน จำกัด, 2543:740 หน้า.
3. Singh K. (Editor). Clinical research on *Morinda citrifolia* L. - Noni. Noni Clinical Research Journal 2007;1(1-2):4-19.
4. Hirazumi A, Furusawa E. An immunomodulatory polysaccharide-rich substance from the fruit juice of *Morinda citrifolia* (noni) with antitumour activity. Phytotherapy Research 1999;13:380-7.
5. Furusawa E, Hirazumi A, Story S and Jensen. Antitumour potential of a polysaccharide-rich substance from the fruit juice of *Morinda citrifolia* (noni) on sarcoma 180 ascites tumour in mice. Phytotherapy Research 2003;17:1158-64.
6. Palu AK, Kim AH, West BJ, Deng S, Jensen J, White L. The effects of *Morinda citrifolia* (noni) on the immune system: Its molecular mechanisms of action. Journal of Ethnopharmacology 2008;115:502-6.
7. Li J, Stickel SL, Bouton-Verville H, et al. Fermented noni exudate (fNE) : A mediator between immune system and anti-tumor activity. Oncology Reports 2008;20:1505-9.
8. Wang MY, Su C. Cancer preventive effect of *Morinda citrifolia* (noni). Ann N Y Acad Sci 2001;952:161-8.
9. Stoner GD, Wang L, Seguin C, et al. Multiple berry types prevent N-nitrosomethylbenzylamine-induced esophageal cancer in rats. Pharm Res 2010;27:1138-45.
10. Wong DK. Are immune responses pivotal to cancer patient's long term survival? Two clinical case - study reports on the effects of *Morinda citrifolia* (noni). Hawaii Med J 2004;63(6):182-4.
11. Issell BF, Gotay CC, Pagano I. Research (ISOQOL) Quality of Life (QOL) Assessment in a Phase I Trial of Noni. Journal of the International Society for Quality of Life Research 2005;14(9):P-70/1342.

12. Issell BF, Gotay CC, Pagano I, Franke AA. Using quality of life measures in Phase I clinical trial of noni in patients with advanced cancer to select a Phase II dose. *Journal of Dietary Supplements* 2009;6(4):347-59.
13. Nuksanit T, Rojanapanthu P, Gritsanapan W, et al. Damnacanthal, a noni component, exhibits antitumorigenic activity in human. *Journal of Nutritional Biochemistry* 2012;23:915–923.
14. Muller JC, Bote GGK, Bufalo AC, et al. *Morinda citrifolia* Linn (noni) : In vivo and vitro reproductive toxicology. *Journal of Ethnopharmacology* 2009;121:229-33.
15. Marques NFO, Marques APBM, Iwano AL, et al. Delayed ossification in wistar rats induced by *Morinda citrifolia* L. exposure during pregnancy. *Journal of Ethnopharmacology* 2010;128:85-91.
16. Nandhasri P, Thamaree S, Punjanon T, Kietnum S. Bio-extract concentrated of Thai "Yor" *Morinda citrifolia* effect in analgesic, acute toxicity and human peripheral blood mononuclear cells. *Thammasat Medical Journal* 2011;11(1):8-17.
17. Stadlbauer V, Fickert P, Krisper P, Trauner M, Stauber RE. Hepatotoxicity of noni juice : report of two cases. *World J Gastroenterol* 2005;11(30):4758-60.
18. Mueller BA, Scott MK, Sowinski KM, Prag KA. Noni juice (*Morinda citrifolia*): Hidden potential for hyperkalemia? *American Journal of Kidney Diseases* 2000;35(2):310-2.