

## เอกสารอ้างอิง: เทียนสัตตบุษย์...สมุนไพรในตำรายาไทย

1. Council of Scientific & Industrial Research. The Wealth of India: A dictionary of raw materials and industrial products Vol. VIII. Calcutta: Sree Saraswaty Press Ltd., 1969:394pp.
2. พยอม ตันติวัฒน์. เครื่องเทศ. กรุงเทพฯ: คณะเภสัชศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย, 2527:119 หน้า.
3. ชยันต์ พิเชียรสุนทร แม้นมาส ชวลิต วิเชียร จีรวงศ์. คำอธิบายตำราพระโอสถพระนารายณ์. กรุงเทพมหานคร: สำนักพิมพ์อมรินทร์, 2542.
4. นันทวัน บุญยะประภัศร และคณะ. สมุนไพรไม้พุ่มบ้าน (2). กรุงเทพมหานคร: สำนักพิมพ์ประชาชน, 2541.
5. Ali-Shtayeh MS, Jamous RM, Jamous RM. Plants used during pregnancy, childbirth, postpartum and infant healthcare in Palestine. Complement Ther Clin Pract 2015;21(2):84-93.
6. British Herbal Medicine Association. British Herbal Pharmacopoeia. Bournemouth: Megaron Press, 1983.
7. Bisset NG. Herbal Drugs and Phytopharmaceuticals. London: CRC Press, 1994.
8. Albert-Puleo M. Fennel and anise as estrogenic agents. J Ethnopharmacol 1980;2(4):337-44.
9. Pourgholami MH, Majzoob S, Javadi M, Kamalinejad M, Fanaee GH, Sayyah M. The fruit essential oil of *Pimpinella anisum* exerts anticonvulsant effects in mice. J Ethnopharmacol 1999;66(2):211-5.
10. Boskabady MH, Ramazani-Assari M. Relaxant effect of *Pimpinella anisum* on isolated guinea pig tracheal chains and its possible mechanism(s). J Ethnopharmacol 2001;74(1):83-8.
11. Sahraei H, Ghoshooni H, Hossein Salimi S, Mohseni Astani A, Shafaghi B, Falahi M, et al. The effects of fruit essential oil of the *Pimpinella anisum* on acquisition and expression of morphine induced conditioned place preference in mice. J Ethnopharmacol 2002;80(1):43-7.
12. Singh G, Kapoor IP, Pandey SK, Singh UK, Singh RK. Studies on essential oils: part 10; antibacterial activity of volatile oils of some spices. Phytother Res 2002;16(7):680-2.
13. Ishikawa T, Fujimatu E, Kitajima J. Water-soluble constituents of anise: new glucosides of anethole glycol and its related compounds. Chem Pharm Bull (Tokyo) 2002;50(11):1460-6.
14. Rodrigues VM, Rosa PT, Marques MO, Petenate AJ, Meireles MA. Supercritical extraction of essential oil from aniseed (*Pimpinella anisum* L.) using CO<sub>2</sub>: solubility, kinetics, and composition data. J Agric Food Chem 2003;51(6):1518-23.
15. Kreydiyyeh SI, Usta J, Knio K, Markossian S, Dagher S. Aniseed oil increases glucose absorption and reduces urine output in the rat. Life Sci 2003;74(5):663-73.
16. Tabanca N, Khan SI, Bedir E, Annavarapu S, Willett K, Khan IA, et al. Estrogenic activity of isolated compounds and essential oils of *Pimpinella* species from Turkey, evaluated using a recombinant yeast screen. Planta Med 2004;70(8):728-35.
17. Tabanca N, Bedir E, Ferreira D, Slade D, Wedge DE, Jacob MR. Bioactive constituents from Turkish *Pimpinella* species. Chem Biodivers 2005;2(2):221-32.
18. Kosalec I, Pepeljnjak S, Kustrak D. Antifungal activity of fluid extract and essential oil from anise fruits (*Pimpinella anisum* L., Apiaceae). Acta Pharm 2005;55(4):377-85.

19. Tabanca N, Demirci B, Ozek T, Kirimer N, Baser KH, Bedir E, et al. Gas chromatographic-mass spectrometric analysis of essential oils from *Pimpinella* species gathered from Central and Northern Turkey. *J Chromatogr A* 2006;1117(2):194-205.
20. Tabanca N, Ma G, Pasco DS, Bedir E, Kirimer N, Baser KH, et al. Effect of essential oils and isolated compounds from *Pimpinella* species on NF-kappaB: a target for antiinflammatory therapy. *Phytother Res* 2007;21(8):741-5.
21. Yoshihiro N. Anethole derivatives having inflammatory cytokine production inhibitory action, and food preparations, cosmetics, and antiinflammatory agents containing them. *Jpn Kokai Tokkyo Koho* 2009, 48pp. JP 2009001514 A 20090108.
22. Capek A. Phytoncides of anise and fennel. *Prumysl Potravin* 1956;7:260-2.
23. Gende LB, Maggi MD, Fritz R, Eguaras MJ. Antimicrobial activity of *Pimpinella anisum* and *Foeniculum vulgare* essential oils against *Paenibacillus larvae*. *J Essent Oil Res* 2009;21:91-3.
24. Karnick CR. *Pharmacopoeial Standards of Herbal Plants*. Vol. 1. Delhi: Sri Satguru Publications, 1994.
25. Burkhardt G, Reichling J, Martin R, Becker H. Terpene hydrocarbons in *Pimpinella anisum* L. *Pharm Weekbl Sci* 1986;8(3):190-3.
26. Al-Bayati FA. Synergistic antibacterial activity between *Thymus vulgaris* and *Pimpinella anisum* essential oils and methanol extracts. *J Ethnopharmacol* 2008;116(3):403-6.
27. Akdemir EG. Empirical prediction and validation of antibacterial inhibitory effects of various plant essential oils on common pathogenic bacteria. *Int J Food Microbiol* 2015;202:35-41.
28. Albulushi SMA, Al Saidi H, Amaresh N, Mullaicharam AR. Study of physicochemical properties, antibacterial and GC-MS analysis of essential oil of the aniseed (*Pimpinella anisum* Linn.) in Oman. *From Research & Reviews: J Pharmacogn Phytochem* 2014;2(4):24-33.
29. Al Maofari A, El Hajjaji S, Debbab A, Zaydoun S, Ouaki B, Charof R, et al. Chemical composition and antibacterial properties of essential oils of *Pimpinella anisum* L. growing in Morocco and Yemen. *St Cerc St CICBIA* 2013;14(1):11-6.
30. Kacaniova M, Vukovic N, Horska E, Salamon I, Bobkova A, Hleba L, et al. Antibacterial activity against *Clostridium* genus and antiradical activity of the essential oils from different origin. *J Environ Sci Health B* 2014;49(7):505-12.
31. Shukla NK, Shankar U. Antibacterial activity of carom seed/thyme (*Trachyspermum ammi*), anise (*Pimpinella anisum*) and dry ginger (*Zingiber officinal*) against *V. cholera*, *V. parahaemolyticus* and *V. vulnificus*. *WJPPS* 2016;5(3):1621-31.
32. Mahady GB, Pendland SL, Stoia A, Hamill FA, Fabricant D, Dietz BM, et al. *In vitro* susceptibility of *Helicobacter pylori* to botanical extracts used traditionally for the treatment of gastrointestinal disorders. *Phytother Res* 2005;19(11):988-91.
33. Robles-Zepeda RE, Velazquez-Contreras CA, Garibay-Escobar A, Galvez-Ruiz JC, Ruiz-Bustos E. Antimicrobial activity of Northwestern Mexican plants against *Helicobacter pylori*. *J Med Food* 2011;14(10):1280-3.

34. Al Salhi M, Devanesan S, Alfuraydi AA, Vishnubalaji R, Munusamy MA, Murugan K, et al. Green synthesis of silver nanoparticles using *Pimpinella anisum* seeds: antimicrobial activity and cytotoxicity on human neonatal skin stromal cells and colon cancer cells. *Int J Nanomedicine* 2016;11:4439-49.
35. Sharma A, Patel VK, Rawat S, Ramteke P, Verma R. Identification of the antibacterial component of some Indian medicinal plants against *Klebsiella pneumoniae*. *Int J Pharm Pharm Sci* 2010;2(3):123-7.
36. Felsociova S, Kacaniova M, Horska E, Hleba L, Petrova J, Rovna K, et al. Antifungal activity of essential oils against selected terverticillate penicillia. *Ann Agric Environ Med* 2015;22(1):38-42.
37. Naeini A, Naseri M, Kamalinejad M, Khoshzaban F, Rajabian T, Nami H, et al. Study on anti-*Candida* effects of essential oil and extracts of Iranian medicinal plants, *in vitro*. *Fasnamah-i Giyahan-i Daruyi* 2011;10(38):163.
38. Lee J-B, Yamagishi C, Hayashi K, Hayashi T. Antiviral and immunostimulating effects of lignin-carbohydrate-protein complexes from *Pimpinella anisum*. *Biosci Biotechnol Biochem* 2011;75(3):459-65.
39. Puertas-Mejia M, Hillebrand S, Stashenko E, Winterhalter P. *In vitro* radical scavenging activity of essential oils from Columbian plants and fractions from oregano (*Origanum vulgare* L.) essential oil. *Flavour Fragr J* 2002;17(5):380-4.
40. Qasem A, Al-Ismael K. Antioxidant activity of alcohol extracts of chamomile flowers, anise seeds and dill seeds in two vegetable oils and two animal fats. *Grasas y Aceites (Sevilla, Spain)* 2004;55(4):345-53.
41. Al-Ismael KM, Aburjai T. Antioxidant activity of water and alcohol extracts of chamomile flowers, anise seeds and dill seeds. *J Sci Food Agric* 2004;84:173-8.
42. Singh G, Kapoor IPS, Singh P, de Heluani CS, Catalan CAN. Chemical composition and antioxidant potential of essential oil and oleoresins from anise seeds (*Pimpinella anisum* L.). *IJEOT* 2008;2(3):122-30.
43. Topal U, Sasaki M, Goto M, Otles S. Chemical compositions and antioxidant properties of essential oils from nine species of Turkish plants obtained by supercritical carbon dioxide extraction and steam distillation. *Int J Food Sci Nutr* 2008;59(7-8):619-34.
44. Nickavar B, Abolhasani FA. Screening of antioxidant properties of seven UMBELLIFERAE fruits from Iran. *Pak J Pharm Sci* 2009;22(1):30-5.
45. Basmacioglu-Malayoglu H, Aktas B, Yesil-Celiktas O. Total phenolic contents and antioxidant activities of the essential oils from some plant species. *Ege Universitesi Ziraat Fakultesi Dergisi* 2011;48(3):211-5.
46. Albayrak S, Aksoy A, Sagdic O, Albayrak S. Antioxidant and antimicrobial activities of different extracts of some medicinal herbs consumed as tea and spices in Turkey. *J Food Biochem* 2012;36(5):547-54.
47. Christova-Bagdassarian VL, Bagdassarian KS, Atanassova MS. Phenolic compounds and antioxidant capacity in Bulgarian plans (dry seeds). *Int J Adv Res* 2013;1(9):186-97.
48. Womeni HM, Tonfack Djikeng F, Tiencheu B, Linder M. Antioxidant potential of methanolic extracts and powders of some Cameroonian spices during accelerated storage of soybean oil. *Adv Biol Chem* 2013;3(3):304-13.

49. Lugemwa FN, Snyder AL, Shaikh K. Determination of radical scavenging activity and total phenols of wine and spices: a randomized study. *Antioxidants* 2013;2(3):110-21.
50. Zielinski AAF, Haminiuk CWI, Alberti A, Nogueira A, Demiate IM, Granato D. A comparative study of the phenolic compounds and the *in vitro* antioxidant activity of different Brazilian teas using multivariate statistical techniques. *Food Res Int* 2014;60:246-54.
51. Ziyatdinova GK, Budnikov HC. Evaluation of the antioxidant properties of spices by cyclic voltammetry. *J Anal Chem* 2014;69(10):990-7.
52. Martins N, Barros L, Santos-Buelga C, Ferreira ICFR. Antioxidant potential of two Apiaceae plant extracts: A comparative study focused on the phenolic composition. *Ind Crops Prod* 2016;79:188-94.
53. Iannarellia R, Marinellia O, Morellib MB, Santonia G, Amantinic C, Nabissia M, et al. Aniseed (*Pimpinella anisum* L.) essential oil reduces pro-inflammatory cytokines and stimulates mucus secretion in primary airway bronchial and tracheal epithelial cell lines. *Ind Crops Prod* 2018;114:81-6.
54. Elsaid FG. Diabetes mellitus and some Egyptian's volatile oils. *Egypt J Hosp Med* 2008; 30:39-50.
55. Al Salhi M, Devanesan S, Alfuraydi AA, Vishnubalaji R, Munusamy MA, Murugan K, et al. Green synthesis of silver nanoparticles using *Pimpinella anisum* seeds: antimicrobial activity and cytotoxicity on human neonatal skin stromal cells and colon cancer cells. *Int J Nanomedicine* 2016;11:4439-49.
56. Devanesan S, AlSalhi MS, Vishnubalaji R, Alfuraydi AA, Alajez NM, Alfayez M, et al. Rapid biological synthesis of silver nanoparticles using plant seed extracts and their cytotoxicity on colorectal cancer cell lines. *J Clust Sci* 2017;28(1):595-605.
57. Karimzadeh F, Hosseini M, Mangeng D, Alavi H, Hassanzadeh GR, Bayat M, et al. Anticonvulsant and neuroprotective effects of *Pimpinella anisum* in rat brain. *BMC Complement Altern Med* 2012;12:76.
58. Janahmadi M, Farajnia S, Vatanparast J, Abbasipour H, Kamalinejad M. The fruit essential oil of *Pimpinella anisum* L. (UMBELLIFERAE) induces neuronal hyperexcitability in snail partly through attenuation of after-hyperpolarization. *J Ethnopharmacol* 2008;120(3):360-5.
59. Abdul-Hamid M, Gallaly SR. Ameliorative effect of *Pimpinella anisum* oil on immunohistochemical and ultrastructural changes of cerebellum of albino rats induced by aspartame. *Ultrastruct Pathol* 2014;38(3):224-36.
60. Ishigaki Y, Tanaka H, Akama H, Ogara T, Uwai K, Tokuraku K. A microliter-scale high-throughput screening system with quantum-dot nanoprobe for amyloid- $\beta$  aggregation inhibitors. *PLoS One* 2013;8(8):e72992.
61. Gamberini MT, Rodrigues DS, Rodrigues D, Pontes VB. Effects of the aqueous extract of *Pimpinella anisum* L. seeds on exploratory activity and emotional behavior in rats using the open field and elevated plus maze tests. *J Ethnopharmacol* 2015;168:45-9.
62. Aboelnaga SMH. Effect of some levels of cardamom, clove and anise on hepatotoxicity in rats caused by CCl<sub>4</sub>. *WASJ* 2015;33(6):854-65.
63. Cengiz N, Özbek H, Him A. Hepatoprotective effects of *Pimpinella anisum* seed extract in rats. *Pharmacologyonline* 2008;3:870-4.

64. El-Sayed MGA, Elkomy A, Samer S, ElBanna AH. Hepatoprotective effect of *Pimpinella anisum* and *Foeniculum vulgare* against carbon tetrachloride induced fibrosis in rats. *WJPPS* 2015;4(6):78-88.
65. Jamshidzadeh A, Heidari R, Razmjou M, Karimi F, Shayesteh MRH, Moein MR, et al. An *in vivo* and *in vitro* investigation on hepatoprotective effects of *Pimpinella anisum* seed essential oil and extracts against carbon tetrachloride-induced toxicity. *Iran J Basic Med Sci* 2015;18(2):205-11.
66. Asadollahpour A, Abdollahi M, Rahimi R, Rahimi R. *Pimpinella anisum* L. fruit: Chemical composition and effect on rat model of nonalcoholic fatty liver disease. *J Res Med Sci* 2017;22:37.
67. Amina B, Nadia AH, Kahloulakhaled, Nesrine S, Abdelkader A. Nephroprotective effect of *Pimpinella anisum* L. aqueous extract against lead toxicity: *In vivo* study. *IJGP* 2016;10(2):86-91.
68. Al Mofleh IA, Alhaider AA, Mossa JS, Al-Soohaibani MO, Rafatullah S. Aqueous suspension of anise "*Pimpinella anisum*" protects rats against chemically induced gastric ulcers. *World J Gastroenterol* 2007;13(7):1112-8.
69. Sharaf A, Goma N. Phytoestrogens and their antagonism to progesterone and testosterone. *J Endocrinol* 1965;31:289-90.
70. Kassi E, Papoutsis Z, Fokialakis N, Messari I, Mitakou S, Moutsatsou P. Greek plant extracts exhibit selective estrogen receptor modulator (SERM)-like properties. *J Agric Food Chem* 2004;52(23):6956-61.
71. Hosseinzadeh H, Tafaghodi M, Abedzadeh S, Taghiabadi E. Effect of aqueous and ethanolic extracts of *Pimpinella anisum* L. seeds on milk production in rats. *J Acupunct Meridian Stud* 2014;7(4):211-6.
72. Javan R, Javadi B, Feyzabadi Z. Breastfeeding: A review of its physiology and galactagogue plants in view of traditional Persian medicine. *Breastfeed Med* 2017;12(7):401-9.
73. Foong SC, Tan ML, Marasco LA, Ho JJ, Foong WC. Oral galactagogues for increasing breast-milk production in mothers of nonhospitalised term infants. *Cochrane Libr* 2015;4:CD011505
74. Shahamat Z, Abbasi-Maleki S, Mohammadi MS. Evaluation of antidepressant-like effects of aqueous and ethanolic extracts of *Pimpinella anisum* fruit in mice. *Avicenna J Phytomed* 2016;6(3):322-8.
75. Shaalan EA, Canyon DV. Inconsequential study on larvicidal efficacy of anise and celery seed extracts indicates that standards in bioinsecticide screening are necessary. *Curr Bioact Compd* 2008;4(1):33 - 5.
76. Pavela R. Insecticidal properties of *Pimpinella anisum* essential oils against the *Culex quinquefasciatus* and the non-target organism *Daphnia magna*. *J Asia Pac Entomol* 2014;17(3):287-93.
77. Kimbaris AC, Koliopoulos G, Michaelakis A, Konstantopoulou MA. Bioactivity of *Dianthus caryophyllus*, *Lepidium sativum*, *Pimpinella anisum*, and *Illicium verum* essential oils and their major components against the West Nile vector *Culex pipiens*. *Parasitol Res* 2012;111(6):2403-10.
78. Khater HF, Hanafy A, Abdel-Mageed AD, Ramadan MY, El-Madawy RS. Control of the myiasis-producing fly, *Lucilia sericata*, with Egyptian essential oils. *Int J Dermatol* 2011;50(2):187-94.
79. Koriem KMM, Arbid MS, El-Gendy NF. The protective role of anise oil in oxidative stress and genotoxicity produced in favism. *J Diet Suppl* 2016;13(5):505-21.
80. Ghazanfarpour M, Sadeghi R, Abdollahian S, Latifnejad RR. The efficacy of Iranian herbal medicines in alleviating hot flashes: A systematic review. *Int J Reprod Biomed (Yazd)* 2016;14(3):155-66.

81. Nahidi F, Kariman N, Simbar M, Mojab F. The study on the effects of *Pimpinella anisum* on relief and recurrence of menopausal hot flashes. *Iran J Pharm Res* 2012;11(4):1079-85.
82. Ghoshegir SA, Mazaheri M, Ghannadi A, Tanhaee M, Feizi A, Babaeian M, et al. *Pimpinella anisum* in modifying the quality of life in patients with functional dyspepsia: A double-blind randomized clinical trial. *J Res Med Sci* 2014;19(12):1118-23.
83. Ghoshegir SA, Mazaheri M, Ghannadi A, Tanhaee M, Feizi A, Babaeian M, et al. *Pimpinella anisum* in the treatment of functional dyspepsia: A double-blind, randomized clinical trial. *J Res Med Sci* 2015;20(1):13-21.
84. Abd El-Salam M, Ammar NM, Yassin N, Ezzeldin N, Zikri EN, El-Anssary AK, et al. A clinico-pharmacological assessment of a herbal preparation for the treatment of bronchial asthma. *WJMS* 2015;12(2):115-24.
85. Vazifekah S, Shams-Ardekani MR, Kamalinejad M, Saderhossein SM, Saedi B, Hosseini H, et al. Evaluation of a novel natural drop for treatment of chronic rhinosinusitis without nasal polyps: a single blind randomized trial. *Int Forum Allergy Rhinol* 2016;6(9):943-9.
86. Mosaffa-Jahromi M, Tamaddon AM, Afsharypuor S, Salehi A, Seradj SH, Pasalar M, et al. Effectiveness of anise oil for treatment of mild to moderate depression in patients with irritable bowel syndrome: A randomized active and placebo-controlled clinical trial. *J Evid Based Complementary Altern Med* 2017;22(1):41-6.
87. Ozguven M. Aniseed. *Handbook of herbs and spices, Volume 2*. Cambridge: Woodhead Publishing Limited; 2012. 138-50.
88. Samojlik I, Mijatovic V, Petkovic S, Skrbic B, Bozin B. The influence of essential oil of aniseed (*Pimpinella anisum* L.) on drug effects on the central nervous system. *Fitoterapia* 2012;83(8):1466-73.
89. Zablocka-Slowinska K, Grajeta H, Jawna K, Biernat J. Interactions between preparations containing female sex hormones and dietary supplements. *Adv Clin Exp Med* 2014;23(4):657-63.
90. Committee on Herbal Medicinal Products (HMPC). Assessment report on *Pimpinella anisum* L., fructus and *Pimpinella anisum* L., aetheroleum. [Internet]. 2012 [cited 2018 Feb 23]. Available from: [http://www.ema.europa.eu/docs/en\\_GB/document\\_library/Herbal\\_-\\_HMPC\\_assessment\\_report/2014/06/WC500168971.pdf](http://www.ema.europa.eu/docs/en_GB/document_library/Herbal_-_HMPC_assessment_report/2014/06/WC500168971.pdf)
91. Samojlik I, Petkovic S, Stilinovic N, Vukmirovic S, Mijatovic V, Bozin B. Pharmacokinetic herb-drug interaction between essential oil of aniseed (*Pimpinella anisum* L., Apiaceae) and acetaminophen and caffeine: a potential risk for clinical practice. *Phytother Res* 2016;30(2):253-9.
92. Swamy PL, Prasad VGNV, Govind V, Rao TS, Kumar LCA, Rao GS. *In-vitro* investigation on fibrinolytic activity of selected indigenous plants and culinary spices. *Inventi Impact: Ethnopharmacology* 2013;(3):162-5.
93. García-González JJ, Bartolomé-Zavala B, Fernández-Meléndez S, Barceló-Muñoz JM, Miranda Páez A, Carmona-Bueno MJ, et al. Occupational rhinoconjunctivitis and food allergy because of aniseed sensitization. *Ann Allergy Asthma Immunol* 2002;88(5):518-22.

94. Gázquez García V, Gaig Jané P, Bartolomé Zavala B. Aniseed-induced nocturnal tongue angioedema. *J Investig Allergol Clin Immunol* 2007;17(6):406-8.