

## เอกสารอ้างอิง\_คีนัว

1. Grubben GJH, Partohardjono S(eds.). Plant resources of South-East Asia No.10. Cereals. Leiden: Backhuys Publishers, 1996:199 pp.
2. Vega-Gálvez A, Miranda M, Vergara J, Uribe E, Puente L, Martinez EA. Nutrition facts and functional potential of quinoa (*Chenopodium quinoa* Willd.), an ancient Andean grain: A review. *J Sci Food Agric.* 2010;90:2541-7.
3. Gordillo-Bastidas E, Díaz-Rizzolo DA, Roura E, Massanés T, Gomis R. Quinoa (*Chenopodium quinoa* Willd), from nutritional value to potential health benefits: An integrative review. *J Nutr Food Sci.* 2016;6(3):1-10.
4. Jancurová M, Minarovičová L, Dandár A. Quinoa-A review. *Czech J Food Sci.* 2009;27(2):71-9.
5. Navruz-Varli S, Sanlier N. Nutritional and health benefits of quinoa (*Chenopodium quinoa* Willd.). *J Cereal Sci.* 2016;69:371-6.
6. Koziol MJ. Chemical composition and nutritional evaluation of quinoa (*Chenopodium quinoa* Willd.). *J Food Comp Anal.* 1992;5:35-68.
7. Ruales J, de Grijalva Y, Lopez-Jaramillo P, Nair BM. The nutritional quality of an infant food from quinoa and its effect on the plasma level of insulin-like growth factor-1 (IGF-1) in undernourished children. *Int J Food Sci Nutr.* 2002;53(2):143-54.
8. Farinazzi-Machado FMV, Barbalho SM, Oshiiwa M, Goulart R, Pessan Junior, O. Use of cereal bars with quinoa (*Chenopodium quinoa* W.) to reduce risk factors related to cardiovascular diseases. *Cienc Technol Aliment Campinas.* 2012;32(3):239-44.
9. De Carvalho FG, Ovidio PP, Padovan GJ, Jordao Junior AA, Marchini JS, Navarro AM. Metabolic parameters of postmenopausal women after quinoa or corn flakes intake-a prospective and double-blind study. *Int J Food Sci Nutr.* 2014; 65(3):380-5.
10. Zevallos VF, Herencia LI, Chang F, Donnelly S, Ellis HJ, Ciclitira PJ. Gastrointestinal effects of eating quinoa (*Chenopodium quinoa* Willd.) in celiac patients. *Am J Gastroenterol.* 2014;109:270-8.
11. Jenkins DJ, Kendall CW, McKeown-Eyssen G, Josse RG, Silverberg J, Booth GL, et al. Effect of a low-glycemic index or a high-cereal fiber diet on type 2 diabetes: a randomized trial. *JAMA.* 2008;300(23):2742-53.
12. Gabriel SGN, Shakib MCR, Gabriel GN. Effect of pseudocereal-based breakfast meals on the first and second meal glucose tolerance in healthy and diabetic subjects. *Open Access Maced J Med Sci.* 2016;4(4):565-73.
13. Li L, Lietz G, Bal W, Watson A, Morfey B, Seal C. Effects of quinoa (*Chenopodium quinoa* Willd.) consumption on markers of CVD risk. *Nutrients.* 2018;10:1-17.
14. Pasko P, Zagrodzki P, Barton H, Chlopicka J, Gorinstein S. Effect of quinoa seeds (*Chenopodium quinoa*) in diet on some biochemical parameters and essential elements in blood of high fructose-fed rats. *Plant Foods Hum Nutr.* 2010;65:333-8.

15. Graf BL, Poulev A, Kuhn P, Grace MH, Lila MA, Raskin I. Quinoa seeds leach phytoecdysteroids and other compounds with anti-diabetic properties. *Food Chem.* 2014;163:178-85.
16. Foucault AS, Mathé V, Lafont R, Even P, Dioh W, Veillet S, et al. Quinoa extract enriched in 20-hydroxyecdysone protects mice from diet-induced obesity and modulates adipokines expression. *Obesity.* 2011;20:270-7.
17. Meneguetti QA, Brenzan MA, Batista MR, Bazotte RB, Silva DR, Cortez DAG. Biological effects of hydrolyzed quinoa extract from seeds of *Chenopodium quinoa* Willd. *J Med Food.* 2011;14(6):653-7.
18. Takao T, Watanabe N, Yuhara K, Itoh S, Suda S, Tsuruoka Y, et al. Hypocholesterolemic effect of protein isolated from quinoa (*Chenopodium quinoa* Willd.) seeds. *Food Sci Technol Res.* 2005;11(2):161-7.
19. Asao M, Watanabe K. Functional and bioactive properties of quinoa and amaranth. *Food Sci Technol Res.* 2010;16(2):163-8.
20. Dini I, Tenore GC, Dini A. Antioxidant compound contents and antioxidant activity before and after cooking in sweet and bitter *Chenopodium quinoa* seeds. *LWT-Food Sci Technol.* 2010;43:447-51.
21. Pasko P, Barton H, Zagrodzki P, Gorinstein S, Foltá M, Zachwieja Z. Anthocyanins, total polyphenols and antioxidant activity in amaranth and quinoa seeds and sprouts during their growth. *Food Chem.* 2009;115:994-8.
22. Hirose Y, Fujita T, Ishii T, Ueno N. Antioxidative properties and flavonoid composition of *Chenopodium quinoa* seeds cultivated in Japan. *Food Chem.* 2010;119:1300-6.
23. Tang Y, Li X, Chen PX, Zhang B, Hernandez M, Zhang H, et al. Characterisation of fatty acid, carotenoid, tocopherol/tocotrienol compositions and antioxidant activities in seeds of three *Chenopodium quinoa* Willd. genotypes. *Food Chem.* 2015;174:502-8.
24. Tang Y, Li X, Zhang B, Chen PX, Liu R, Tsao R. Characterisation of phenolics, betanins and antioxidant activities in seeds of three *Chenopodium quinoa* Willd. genotypes. *Food Chem.* 2015;166:380-8.
25. Nsimba RY, Kikuzaki H, Konishi Y. Antioxidant activity of various extracts and fractions of *Chenopodium quinoa* and *Amaranthus* spp. seeds. *Food Chem.* 2008;106:760-6.
26. Tang Y, Zhang B, Li X, Chen PX, Zhang H, Liu R, et al. Bound phenolics of quinoa seeds released by acid, alkaline, and enzymatic treatments and their antioxidant and  $\alpha$ -glucosidase and pancreatic lipase inhibitory effects. *J Agric Food Chem.* 2016;64:1712-9.
27. Yao Y, Shi Z, Ren G. Antioxidant and immunoregulatory activity of polysaccharides from quinoa (*Chenopodium quinoa* Willd.). *Int J Mol Sci.* 2014;15:19307-18.
28. Aluko RE, Monu E. Functional and bioactive properties of quinoa seed protein hydrolysates. *J Food Sci.* 2003;68(4):1254-8.

29. Vilcacundo R, Miralles B, Carrillo il W, Hernández-Ledesma B. *In vitro* chemopreventive properties of peptides released from quinoa (*Chenopodium quinoa* Willd.) protein under simulated gastrointestinal digestion. *Food Res Int.* 2018;105:403-11.
30. Graf BL, Cheng DM, Esposito D, Shertel T, Poulev A, Plundrich N, et al. Compounds leached from quinoa seeds inhibit matrix metalloproteinase activity and intracellular reactive oxygen species. *Int J Cosmet Sci.* 2015;37:212-21.
31. Pasko P, Barton H, Zagrodzki P, Izewska A, Krosniak M, Gawlik M, et al. Effect of diet supplemented with quinoa seeds on oxidative status in plasma and selected tissues of high fructose-fed rats. *Plant Foods Hum Nutr.* 2010;65:146-51.
32. Hemalatha P, Bomzan DP, Rao BVS, Sreerama YN. Distribution of phenolic antioxidants in whole and milled fractions of quinoa and their inhibitory effects on  $\alpha$ -amylase and  $\alpha$ -glucosidase activities. *Food Chem.* 2016;199:330-8.
33. Ayyash M, Johnson SK, Liu SQ, Al-Mheiri A, Abushelaibi A. Cytotoxicity, antihypertensive, antidiabetic and antioxidant activities of solid-state fermented lupin, quinoa and wheat by *Bifidobacterium* species: *In vitro* investigations. *LWT-Food Sci Technol.* 2018;95:295-302.
34. Nickel J, Spanier LP, Botelho FT, Gularte MA, Helbig E. Effect of different types of processing on the total phenolic compound content, antioxidant capacity, and saponin content of *Chenopodium quinoa* Willd grains. *Food Chem.* 2016; 209:139-43.
35. Kuljanabhadgavad T, Thongphasuk P, Chamulitrat W, Wink M. Triterpene saponins from *Chenopodium quinoa* Willd. *Phytochemistry.* 2008;69: 1919-26.
36. Gawlik-Dziki U, S´wieca M, Sułkowski M, Dziki D, Baraniak B, Czyz J. Antioxidant and anticancer activities of *Chenopodium quinoa* leaves extracts-*In vitro* study. *Food Chem Toxicol.* 2013;57:154-60.
37. Yao Y, Yang X, Shi Z, Ren G. Anti-inflammatory activity of saponins from quinoa (*Chenopodium quinoa* Willd.) seeds in lipopolysaccharide-stimulated RAW 264.7 macrophages cells. *J Food Sci.* 2014;79(5):H1018-H1023.
38. Woldemichael GM, Wink M. Identification and biological activities of triterpenoid saponins from *Chenopodium quinoa*. *J Agric Food Chem.* 2001;49:2327-32.
39. Pagno CH, Costa TMH, De Menezes EW, Benvenuti EV, Hertz PF, Matte CR. Development of active biofilms of quinoa (*Chenopodium quinoa* W.) starch containing gold nanoparticles and evaluation of antimicrobial activity. *Food Chem.* 2015;173:755-62.
40. Verza SG, Silveira F, Cibulski S, Kaiser S, Ferreira F, Gosmann G, et al. Immunoadjuvant activity, toxicity assays, and determination by UPLC/QTOF-MS of triterpenic saponins from *Chenopodium quinoa* seeds. *J Agric Food Chem.* 2012;60:3113-8.
41. Hong J, Convers K, Reeves N, Temprano J. Anaphylaxis to quinoa. *Ann Allerg Asthma Immunol.* 2013;110(1):60-1.

42. Astier C, Moneret-Vautrin DA, Puillandre E, Bihain BE. First case report of anaphylaxis to quinoa, a novel food in France. *Allergy* 2009;64:819-20.
43. Lopez DEQ, Zavala BB, Ortiz I. Cross-reactivity between buckwheat and quinoa in a patient with eosinophilic esophagitis caused by wheat. *J Investig Allergol Clin Immunol.* 2014;24(1):56-71.