

เอกสารอ้างอิงของมะเขือเทศ 17/3

1. Steinegger E, Haensel R. *Pharmakognosie*.5. Berlin: Springer-Verlag,1992.
2. Nuhn P. *Naturstoffchemie*.3. Stuttgart: S. Hirzel Verlag, 1997.
3. Teuscher E, Lindequist U. *Biogene Gifte*. Stuttgart Gustav Fischer, 1994.
4. Konowalchuk J, Speirs JI. Antiviral effect of commercial juices and beverages. *Appl environ microbiol* 1978;35:1219.
5. Shimoi K, Masuda S, Shen B, et al. Radioprotective effects of antioxidative plant flavonoids in mice. *Mutat res* 1996;350(1):153-61.
6. Lim-Sylianco CY, Conchor JA, Jocano AP, et al. Antimutagenic effects of expressions from twelve medicinal plants. *Philippine J Sci* 1986;115(1):23-30.
7. Okamina E, Tsutsumi M, Ozono S, et al. Inhibitory effect of tomato juice on rat urinary bladder carcinogenesis after N-butyl-N-(4-hydroxybutyl) nitrosamine initiation. *Jap J cancer Res (Gann)*. 1998;89(1):22-6.
8. Sakamoto H, Mori H, Olima F, et al. Elevation of serum carotenoids after continual ingestion of tomato juice. *Nippon Siyo, Shokungo Gokkaishi* 1994;47(2):93-9.
9. Piercy KT, Myers RD. Tomato juice, chocolate drink and other fluid suppress volitional drinking of alcohol in the female syrian golden hamster. *Physiol Behav* 1995;57(6):1156-61.
10. Yan xinhaue, et al. Fast acting wrinkle-eliminating liquids. Patent:1,070,105, 1993:4 pp.
11. Hirakoshi J, Ishinabe K, Yatagai K, et al. Urination promoters containing saw-palmetto extracts and lycopene. Patent:Japan Kokai Tokkyo Koho 10 158,182 ,1998:4pp.
12. Francheschi S, Bidoli E, La RC, et al. Tomatoes and risk of digestive-tract cancers. *Int J Cancer* 1994;59(2):181-4.
13. Hirakoshi J, Ishinabe K, Yatagai K, et al. Urination promoters containing saw-palmetto extracts and lycopene. Patent:Japan Kokai Tokkyo Koho 10 158,182, 1998:4pp.
14. Yasukawa K, Yamaguchi A, Arita J, et al. Inhibitory effect of edible plant extracts on 12-O-tetradecanoylphorbol-13-acetate-induced ear oedema in mice. *Phytother Res* 1993;72:185-9.
15. Prochaska HJ, Santamaria AB, Talalay P. Rapid detection on inducers of enzymes that protect against carcinogens. *Proc Nat Acad Sci (USA)* 1992;89:2394-8.
16. Rauscher R, Edenharder R, Platt KL. In vitro antimutagenic and in vivo anticlastogenic effects of carotenoids and solvent extracts from fruits and vegetables rich in carotenoids. *Mutat Res* 1998;413(2):129-42.

17. Maeda H, Katsuki T, Akaike T, et al. High correlation between lipid peroxide radical and tumor-promoter effect: suppression of tumor promotion in the Epstein-barr virus/B-lymphocyte system and scavenging of alkyl peroxide radicals by various vegetable extracts. *Jap J Cancer Res (Gann)* 1992;83(9):923-8.
18. Sekiya K, Fushimi T, Ishikawa N, et al. Regulation of arachidonic acid metabolism in platelets by vegetables. *Biosci Biotech Biochem* 1993;57(4):670-1.
19. Shinohara K, Kuroki S, Miwa M, et al. Antimutagenicity of dialyzates of vegetables and fruits. *Agr Biol Chem* 1988;52(6):1369-75.
20. Osawa T, Ishibashi H, Namiki M, et al. Desmutagenic action of food components on mutagens formed by the sorbic acid nitrite reaction. *Agr Biol Chem* 1978;50(8):1971-7.
21. Murakamia A, Jiwajiinda S, Koshimizu K, et al. Screening for in vitro anti-tumor promoting activities of edible plants from Thailand. *Cancer Lett* 1995;95(1/2):137-46.
22. Pressey R. Invertase inhibitor in tomato fruit. *Phytochemistry* 1994;36(3):543-6
23. Cohen Y, Guegler K, Moesinger E, et al. Fungicidal proteins P14 of tomato; purification and cloning of sequences encoding them. Patent:Pct Int Appl 92 20,800, 1992:36pp.
24. Betz H, Schloesser E. On the use of fungicidal saponins in crop protection. *Tagungsber-Akad Landwirtschaftswiss DDR* 1984;222:179-87.
25. Verhoeff K, Liem JI. Toxicity of tomatine to *Botrytis cinerea* in relation to latency. *Phytopathol Z* 1975;82:333.
26. Weniger B, Rouzier M, Daguilh R, et al. Popular medicine of the central plateau of Haiti.2. Ethnopharmacological inventory. *J Ethnopharmacol* 1986;17(1):13-30.
27. Konji K, Takeshi T, Keiko M, et al. Vasoactive substances in foods. *Nippon Eiyo, Shokuryo Gakkaishi* 1991;44(6):487-92.
28. Melotto E, Greve LC, Labaich JM. Biologically active pectin oligomers in ripening tomato fruits. *Prog Biotechnol* 1996;14:207-17.
29. Hass GM, Ryan CA. Carboxypeptidase inhibitor from ripened tomatoes : purification and properties. *Phytochemistry* 1980;19:1329-33.
30. Giouannucci E, Ascherio A, Rim E, et al. Intake of carotenoids and retinol in relation to risk of prostate cancer. *J Natl Cancer Inst* 1995;87(23):1767-7.
31. Gasperi-campani A, Barbieri L, Battelli MG, et al. On the distribution of ribosome inactivating proteins amongst plants. *J Nat Prod* 1985;48(3):446-54.

32. Malinow M. Cholesterol sequestrant glycosides that inhibit intestinal cholesterol absorption. Patent:PCT Int Appl 95 00, 1995:32 pp.
33. Rao P. Nutrient composition and biological evaluation of defatted tomato (*Lycopersicon esculenta*) seed cake. *Plant Foods Hum Nutr* 1991;41(1):101-6.
34. Wakkary JA, Goodfriend L, Kovacs BA. Isolation and some pharmacological properties of two biologically active substances of crown gall infected tomato plants. *Arch Int Pharmacodyn Ther* 1970;183:289.
35. Celayeta FD. Action of the tissues of various plants on the growth of *Sphacelia segetum*. *Farmacognosia (Madrid)* 1960;20:91-101.
36. Anon. Pharmaceutical and cosmetic compositions containing tomato plant extracts for the treatment of skin diseases. Patent:ISRAELI 78,820, 1987:15pp.
37. Anon. Pharmaceutical and cosmetic compositions containing tomato plant extracts for the treatment of skin diseases. Patent:ISRAELI 78,820, 1987:15pp.
38. Itob E, Kim CS, Horike M. A feeding deterrent for thrips palmi karny (*Thysanoptera*:Thripidae) found in tomato leaves. *Nippon Oyo Dobutsu Konchu Gakkaishi* 1994;38(2):109-20.
39. Singh J, Dubey AK, Triparhi NN. Antifungal activity of *Mentha spicata*. *Int J Pharmacog* 1994;32(4):314-9.
40. Singh KV, Pathak RK. Effect of leaves extracts of some higher plants on spore germination of *Ustilago maydes* and *U. nuda*. *Fitoterapia* 1984;55(5):318-20.
41. Kone-bamba D, Pelissier Y, Onoukou ZF, et al. Hemostatic activity of 216 plants used in traditional medicine in the Ivory coast. *Plant Med Phytother* 1987;21(2):122-30.
42. Medina FR, Woodbury R. Terrestrial plants molluscicidal to lymnaeid hosts of *Fascioliasis hepatica* in Puerto Rico. *J Agr Univ Puerto* 1979;63:366-76.
43. Hildebrand DF, Brown GC, Jackson DM, et al. Effects of some leaf-emitted volume compounds on aphid popular increase. *J Chem Ecol* 1993;19(9):1875-87.
44. Hirono C, Yasumi K, Itoh E, et al. Feeding deterrent for thrips palmi leray (*Thysanoptera* : Thripidae) found in tomato leaves. Isolation and identification. *Nippon Oyo Dobutsu Koncha Gakkaishi* 1994;38(2):109-20.
45. Lin, SYH, Taumble JT, Kumamoto J. Activity of volatile compounds in glandular trichomes of *Lycopersicon* species against two insect herbivores. *J Chem Ecol* 1987;13(4):837-50.

46. Duttey SS, Isman MB. Inhibition of insect larval growth by phenolics in glandular trichomes of tomato leaves. *Experientia* 1988;52(6):1359-75.
47. John SL, et al. 2,3-Dihydrofernesoic acid, a unique terpene from trichems of *Lycopersicon hirsuta*, repels spider mites. *J Chem Ecol* 1993;19(12):1981-97.
48. Venture MU, Vandermin JD. Toxicity of 2-tridecanone and 2-undecanone allelochemicals present in tomato (*Lycopersion* spp.) for *Phthorimaea operculella* (Zell) larvae. *Sci Agric* 1995;52(3):453-61.
49. Hammerschlag F, Mace ME. Antifungal activity of extracts from *Fusarium* wilt-susceptible and-resistant tomato plants. *Phytopathology* 1975;65:93.
50. Carnes MG, Brenner ML, Andersen CR. Comparison of reversed-phase high-pressure liquid chromatography with sephadex LH-20 for cytokinin analysis of tomato root pressure exudate. *J chromatogr* 1975;108:95.
51. Nagaoka T, Ohra J, Yoshihara T, et al. Fungitoxic compound for the roots of tomato stock. *Nippon Shokubutsu Byori Gakkhaiho* 1995;61(2):103-8.
52. Aswal BS, Bhakuni DS, Goel AK, et al. Screening of Indian plants for biological activity:Part X. *Indian J Exp Biol* 1984;22(6):312-32.
53. Roychoudhury R. Effect of extracts of certain *Solanaceous* plants on plant virus infection. *Acta Bot Indica* 1980;81:91-4.
54. อัญชลี ตันท์ศุภศิริ และคณะ. การใช้สารสกัดอินสีกัดจากมะเขือเทศสีดา. *สารศิริราช* 2534;43(12):918-25.
55. *Pharmaceutical Research* 1997;14(5).
56. Naisbett B, Woodley J. The potential use of tomato lectin for oral drug delivery:2. Mechanism of uptake in vitro. *Int J Pharm* 1994;110(2):127-36.
57. Stahl W, Sies H. Uptake of lycopene and its geometrical isomers is greater from heat-processed than from unprocessed tomato juice in haman. *J Nutr* 1992;122(11):2161-6.
58. Sharoni Y, Levy J. Lycopene the major tomato carotenoid inhibits endometral and lung cancer cell growth. *Proc J Proc Int Cancer Congr, Free Pap Posters, 16th, 1994;1:641-5.*
59. Altonassova V, Pevitcharova G. Study of the influence of food products on serum nitrite content and rat mortality. *Med Nutri* 1997;33(5):199-203.
60. Formulation containing carotenoids and procarotenoids combinedwith polyphenols for the prevention of the damages due to an abnormal production of free redicals. Patent:US US 5,648,377, 1997:6pp.

61. Yoau S, Eva G, Moshe R, et al. Effects of lycopene-enriched tomato oleoresin on 7,12-dimethylbenz[a]anthracene-induced rat memmory tumors. *Cancer Detect Prev* 1997;21(2):118-23.
62. Spinu K, et al. Antiviral activity of tomatoside from *Lycopersicon esculentum* Mill. *Adv Exp Med Biol* 1996;404:505-9.

เอกสารอ้างอิงของหม่อน

12/2

1. วุฒิ วุฒิธรรมเวช. เกษตรกรรมไทยรวมสมุนไพร ฉบับปรับปรุงใหม่. พิมพ์ครั้งที่ 2. กรุงเทพฯ:โอ.เอส.พรินติ้งเฮ้าส์, 2540.
2. Naito K, Hamamura Y. Trace constituent in mulberry leaves III. Isolation of β -sitosterol from mulberry leaves. Nippon Nogeikagaku Kaishi 1961;35:848-50.
3. Goto M, Kamada M, Imai S, et al. Mulberry and related materials I. Takeda Kenkyusho Nempo 1965;24:55-65.
4. Alkofahi A, Al-Khalil S. Studies on the chemistry of the leaves of *Morus alba*. Bull Fac Pharm (Cairo Univ.) 1996;34(1):59-62.
5. Yamazaki M, Nkamura N, Kurioka A. Identification of isofurosterol in mulberry leaves, *Morus alba*. Nippon Sanshigaku Zasshi 1997;66(2):136-7.
6. Ogawa S, Nishimoto S, Hirayama H, Taniguchi S, Takemoto T. Extraction of inokosterone from mulberry. Patent:Japan 70 36,504, 1970:2 pp.
7. Chen F, Nakashima N, Kimura I, Kimura M, Asano N, Koya S. Potentiating effects on pilocarpine-induced saliva secretion, by extracts and N-containing sugars derived from mulberry leaves, in streptozotocin-diabetic mice. Biol Pharm Bull 1995;18(12):1676-80.
8. Talyshinskii GM. Formation of rutin in leaves of mulberry trees. Dokl Akad Nauk Azerb SSR 1967;23(1):63-6.
9. Naito K. Trace components in mulberry leaves. VII. Isolation of rutin and quercetin from mulberry leaves. Nippon Nogeikagaku Kaishi 1968;42(7):423-5.
10. Naito K. Trace constituents of mulberry leaves. VIII. Isolation of quercetin glycoside from mulberry leaves. Nippon Nogeikagaku Kaishi 1968;42(8):450-3.
11. Onogi A, Osawa K, Yasuda H, Sakai A, Morita H, Itokawa H. Flavonol glycosides from the leaves of *Morus alba* L. Shoyakugaku Zasshi 1993;47(4):423-5
12. Takasugi M, Naguo S, Masamune T, Shirata A, Takahashi K. Studies on phytoalexins of the Moraceae, *U. chalconoracin*, a natural Diels-Alder Adduct from diseased mulberry. Chem Lett 1980:1573-6.
13. El-Khrisy F, Nassar MI, Abu-Mustafa EA. Constituents of *Morus alba* leave. Fitoterapia 1992;63(1):92.
14. Matsuyama S, Kuwahara Y, Suzuki T. A new 2-arylbenzofuran, omega-hydroxy maracin N from mulberry leaves. Agr Biol Chem 1991;55(5):1409-10.
15. Ferent M. Ceramide extraction process. Patent:Fr Demande Fr 2,739,853, 1997:11 pp.
16. Dudkin MS, Shkantova NG. Chemical composition of *Morus alba* leave. Khim Drev 1982;1:101-2.

17. Jayal MM, Kehar ND. Nutritive value of mulberry tree leaves. J Dairy Sci 1962;15:21-7.
18. Talyshinskii GM. Carotene and vitamin C levels in the leaves of mulberry trees. Tr Inst Genet Selek Akad Nauk Azerb SSR 1970;6:264-7.
19. Kirimura J, Suzuki T and Hashimoto Y. The vitamin-B complex content of mulberry leaves. Sanshi Shikensho Hokoku 1962;17:545-57.
20. Iwanari Y. Detection of volatile neutral components in mulberry leaves harvested in late autumn. Nippon Sanshigaku Zasshi 1973;42(5):403-5.
21. Qiu F, Komatsu K, Kawasaki K, Saito K, Yao X, Kano Y. A novel stilbene glucoside, oxyresveratrol 3'-O- β -glucopyranoside from the root bark of *Morus alba*. Planta Med 1996;62:559-61.
22. Hano Y, Tsubura H, Nomura T. Constituents of the cultivated mulberry tree XXXVIII. Structures of kuwanons Y and Z, two new stilbene derivative from *Morus alba* L. Heterocycle 1986;24(9):2603-10.
23. Luo SD, Nemeč J, Ning BM. Anti-HIV flavonoids from *Morus alba*. Yun-nan Chih Wu Yen Chiu 1995;17(1):89-95.
24. Nomura T, Fukai T. Kuwanon G, a new flavone derivative from the root barks of the cultivated mulberry tree (*Morus alba*). Chem Pharm Bull 1980;28:2548-52.
25. Nomura T, Fukai T, Narita T. Hypotensive constituent, kuwanon H, a new flavone derivative from the root bark of the cultivated mulberry tree (*Morus alba*). Heterocycles 1980;14:1943-51.
26. Nomura T, Fukai T, Hamo Y, et al. Structures of natural Diels-Alder adducts of *Morus* root barks. Tennen Yuki Kagobutsu Toronkai Koen Yoshishu 1981;24:151-8.
27. Zenyaku Kogyo Co, Ltd. Sanggenone B from mulberry roots. Patent:JPN Kokai Tokkyo Koho JP 82,146,718, 1982:5 pp.
28. Nomura T, Fukai T, Uno J, Arai T. Mulberrofuran A, a new isoprenoid 2-arylbenzofuran from the root bark of the cultivated mulberry tree (*Morus alba* L.). Heterocycles 1978;9(11):1593-601.
29. Nomura T, Fukai T. Mulberrofuran B, a new isoprenoid 2-arylbenzofuran from the root bark of cultivated mulberry tree. Planta Med 1981;42(2):197-9.
30. Rao AV, Deshpande VH, Shastri RK, Tavale SS, Dhaneshwar NN. Structures of albanols A and B, two novel phenols from *Morus alba* bark. Tetrahedron Lett. 1983;24(29):3013-6.
31. Deshpande VH, Srinivasan R, Rama Ras AV. Wood phenolics of *Morus* species. IV. Phenolics of the heartwood of five *Morus* species. Indian J Chem 1975;13(5):453-7.
32. Deshpande VH, Wakharkar PV, Rao AVR. Wood phenolics of *Morus* species : Part V. Isolation of a new flavone, mulberranol and a novel phenol, alboctalol from *Morus alba*. Indian J Chem Sect B 1976;14B(9):647-50.

33. Kasyanov GI, Pekhov AV, Shaftan EA, Klimova ES, Shishkov GZ. Carbon dioxide extract of *Morus alba* L. fruit. *Rastit Resur* 1976;12(4):597-9.
34. Yang CH, Tsai TC. Anthocyanins in mulberry fruit. *Shipin Kexue* 1994;21(5):319-30.
35. Shapyan GG, Minasyan SM, Astabatsyan GA, et al. Study of biochemical indexes and technological properties of the mulberry. *Konservn Ovoshchesush Prom-st* 1981;6:35-6.
36. Makhmudbekova NI. Biochemical study of fruits of mulberry trees grown on the Apsheron Peninsula. *Doklady Akad Nauk Azerbaidshan SSR* 1961;17:621-6.
37. Ono K, Nakane H, Meng ZM, Ose Y, Sakai Y, Mizuno M. Differential inhibitory effects of various herb extracts on the activities of reverse transcriptase and various DNA polymerases. *Chem Pharm Bull* 1989;37(7):1810-2.
38. Luo S, Nemeč J, Ning B. Anti-HIV flavonoids from *Morus alba*. *Yannan Zhiwu Yanjiu* 1995;17(1):89-95.
39. Luo S, Ning B. HIV-1 inhibitors for treatment of AIDS. Patent:Faming Zhuanli Shenqing Gongkai Shuomingshu CN 1,107,152, 1995:11 pp.
40. Chen FJ, Nakanishima N, Kimura M. Hypoglycemic activity and mechanism of extracts from mulberry leaves and cortex mori radices in streptozotocin-induced diabetic mice. *Yakugaku Zasshi* 1995;115(6):476-82.
41. Sharaf A, Mansour MY. Pharmacological studies on the leaves of *Morus alba*, with special reference to its hypoglycemic activity. *Panta Med* 1964;12(1):71-6.
42. Yamada H, Oya I, Nagai T, Matsumoto T, Kiyohara H, Omura S. Screening of α -glucosidase II inhibitors from Chinese herbs and its application on the quality control of mulberry bark. *Shoyakugaku Zasshi* 1993;47(1):47-55.
43. Sanavova MK, Rakhimov DA. Plant polysaccharides VII. Polysaccharides of *Morus* and their hypoglycemic activity. *Chem Nat Compd* 1997;33(6):617-9.
44. Onogi A, Osawa K, Yasuda H, Sakai A, Morita H, Itokawa H. Flavonoid glycoside from the leave of *Morus alba* L. *Shoyakugaku Zasshi* 1993;47(4):423-5.
45. Ionescu TC, Popa E, Mirodon Z, Simionescu M, Mineu I. Effect of a mixture of plants on the metabolic equilibrium in patients with diabetes mellitus of the 2nd type. *Rev Med Interna* 1989;41(2):185-92.
46. Yamatake Y, Shibata M, Nagai M. Pharmacological studies on root bark of mulberry tree (*Morus alba* L.) *Jap J Pharmacol* 1976;26:461.
47. Zenyaku Kogyo Co,ltd. 8-Cyclohexynylflavones from *Mulberry* species. Patent: JPN Kokai Tokkyo Koho JP 81,123,979, 1981.

48. Nomura T, Fukai T, Hano Y, Uzawa J. Structure of sanggenon C, a natural hypotensive Diels-Alder adduct from Chinese crude drug 'Sang Bai-Pi' (*Morus* root barks). *Heterocycles* 1981;16(12):2141-8.
49. Zenyaku Kogyo Co, Ltd. Sanggenone D. Patent:JPN Kokai Tokkyo Koho JP 58 41,894 ,1983.
50. Oshima Y, Konno C, Hikino H. Validity of oriental medicine. Part 28. Structure of moracenin D, a hypotensive principle of *Morus* root bark. *Heterocycles* 1981;16(6):979-82.
51. Fukai T, Hano Y, Hirakura K, Nomura T, Uzawa J, Fukushima K. Structures of mulberrofurans F and G, two natural hypotensive Diels-Alder type adducts from the cultivated mulberry tree. *Heterocycles* 1984;22(3):473-7.
52. Lee HK, Chung YS. Antifungal activities of medicinal plants in Korea. *Kisul Yongerso Pogo* 1963;2:76-8.
53. Winter AG, Willeke L. Investigations of plant antibiotics IV. Inhibitors in plant foliage. *Naturwissenschaften* 1952;39:45.
54. Zenyaku Kogyo Co,ltd. Isolation of microbicidal kuwanone L. Patent:JPN Kokai Tokkyo Koho JP 58,150,584 ,1983.
55. Nomura T, Fukai T, Uno J, Arai T. Mulberrofuran A, a new isoprenoid 2-arylbenzofuran from the root bark of cultivated mulberry tree. *Heterocycles* 1478;9:1593-600.
56. Park WJ, Lee HJ, Yang SG. The inhibitory effect of sanggenon C from the root bark of *Morus alba* L. on the growth and cellular adherence of *Streptococcus mutans*. *Yakhak Hoechi* 1990;34(6):434-8.
57. Yen GC, Wu SC, Du PD. Extraction and identification of antioxidants from the leaves of mulberry (*Morus alba* L.). *J Agr Food Chem*
58. Huang DQ, Xiao H, Fu WQ, Zhao B, Xin WJ. Effects of extracts of mulberry leaves (*Morus alba* L.) on scavenging oxygen free radicals in vitro and on the life span of *Drosophila melanogaster*. *Fujian Shifan Daxue Xuebao Ziran Kexueban* 1995;11(1):85-90.
59. Shin NH, Lee KS, Kang SH, Min KR, Lee SH, Kim VS. Inhibitory effects of herbal extracts on Dopa oxidase activity of tyrosinase. *Nat Prod Sci* 1997;3(2):111-121.