

เอกสารอ้างอิง (อาหารไทยกับการป้องกันมะเร็ง)

1. Itokawa H, Morita H, Sumitomo T, Totsuka N, Takeya K. Antitumour principles from *Alpinia galanga*. *Planta Med* 1987;53(1):32-3.
2. Murakami A, Ohigashi H, Koshimizu K. Possible anti-tumour promoting properties of traditional Thai food items and some of their active constituents. *Asia Pacific J Clin Nutr* 1994;3:185-91.
3. Tiwawech D, Hirose M, Futakuchi M, Lin C, Thamavit W, Ito N, Shirai T. Enhancing effects of The edible plants on 2-amino-3, 8-dimethylimidazo (4,5-f) quinoxaline-hepatocarcinogenesis in a rat medium-term bioassay. *Cancer Lett* 2000;158(2):195-201.
4. Tanaka T, Makita H, Kawamori T, *et al.* A xanthine oxidase inhibitor 1'-acetoxychavicol acetate inhibits azoxymethane-induced colonic aberrant crypt foci in rats. *Carcinogenesis* 1997;18(5):1113-8.
5. Tanaka T, Kawabata K, Kakumoto M, *et al.* Chemprevention of azoxymethane-induced rat colon carcinogenesis by a xanthine oxidase inhibitor, 1'-acetoxychavicol acetate. *Jpn J Cancer Res* 1997;88(9):821-30.
6. Yozo K, Dai N, Hiroyuki A, *et al.* Prevention by 1'-acetoxychavicol acetate of the induction but not growth of putative preneoplastic, glutathione S-transferase placental form-positive, focal lesions in the livers of rats fed a choline-deficient, L-amino acid-defined diet. *Carcinogenesis* 1998;19(10):1809-14.
7. Moffatt J, Hashimoto M, Kojima A, *et al.* Apoptosis induced by 1'-acetoxychavicol acetate in Ehrlich ascites tumor cells is associated with modulation of polyamine metabolism and caspase-3 activation. *Carcinogenesis* 2000;21(12):2151-7.
8. Murakami A, Toyota K, Ohura S, Koshimizu K, Ohigashi H. Structure-activity relationships of (1'-S)-1'-acetoxychavicol acetate, a major constituent of a southeast Asian condiment plant *Languas galanga*, on the inhibition of tumor-promoter-induced epstein-barr virus activation. *J Agric Food Chem* 2000;48:1518-23.