

## เอกสารอ้างอิง Capsinoids: กลุ่มสารสำคัญในพริกไม่เผ็ด

1. Gannett PM, Nagel DL, Reilly PJ, Lawson T, Sharpe J Toth B. The capsaicinoids: their separation, synthesis, and mutagenicity. J Org Chem 1988;53:1064 - 71.
2. Heresch F, Jurenitsch J. Off-line mass spectrometric monitoring of HPLC effluents - an improved identification and quantitation method for mixtures of similar compounds: natural capsaicinoids. J Chromatographia 1979;12:647 - 50.
3. Sticher O, Soldati F, Joshi RK. Hochleistungsflüssigkeitschromatographische trennung und quantitative bestimmung von capsaicin, dihydrocapsaicin, nordihydrocapsaicin und homodihydrocapsaicin in natürlichen capsaicinoid-gemischen und fructus capsici. J Chromatogr 1987;166:221 - 31.
4. Kopp B, Jurenitsch J. Biosynthese der capsaicinoide in *Capsicum annuum* L. var. *annuum* II. bildung des säurerestes der capsaicinoide aus L-valin, L-leucin und L-isoleucin. Planta Med 1981; 43:272 - 9.
5. Jurenitsch J, Wginger R. Isolierung und struktur von homodihydrocapsaicin II. J. jurenitsch und R. wöginger. Sci Pharm 1982;50:111 - 4.
6. Jurenitsch J, David M, Heresch F, Kubelka W. Nachweis und identifizierung neuer scharfstoffe in *Capsicum*-früchten. Planta Med 1979;36:61 - 7.
7. Bennett DJ, Kirby GW. Constitution and biosynthesis of capsaicin. J Chem Soc (C) 1968:442 - 7.
8. Buck SH, Burks TF. The neuropharmacology of capsaicin: review of some recent observations. J Pharmacol Rev 1986;38:179 - 226.
9. Suzuki T, Iwai K. Constituents of red pepper spices: chemistry, biochemistry, pharmacology and food science of the pungent principle of *Capsicum* species. Chemistry and Pharmacology 1984;23:1984.
10. Yazawa S, Suetome N, Okamoto K, Namiki T. Content of capsaicinoids and capsaicinoid-like substances in fruit of peper (*Capsicum annuum* L.) Hybrids Made with "CH-19 sweet". J Jpn Soc Hortic Sci 1989;58:601 - 7.
11. Watanabe T, Kawada T, Kato T, Harada T, Iwai K. Effect of capsaicin analogs on adrenal catecholamine secretion in rats. Life Sci 1994;54: 369 - 74.
12. Kobata K, Todo T, Yazawa S, Iwai K, Watanabe T. Novel capsaicinoid-like substances, capsiate and dihydrocapsiate, from the fruits of a nonpungent cultivar, CH-19 sweet, of pepper (*Capsicum annuum* L.). J Agric Food Chem 1998;46:1695 - 7.

13. (a) Yazawa S, Tatsuo N, Watanabe T, Fushiki T, Kobata K, Imai M, Setoguchi Y, Hashizume S. Capsaicinoid-Like substances having ester bond. US Patent 6,333,421 B1 December 18, 2001.
13. (b) [www.capsiatenatura.com/glossary.aspx](http://www.capsiatenatura.com/glossary.aspx) 09/11/2010
14. Ohnuki K, Niwa S, Maeda S, Inoue N, Yazawa S, Fushiki T. CH-19 sweet, a nonpungent cultivar of red pepper, increased body temperature and oxygen consumption in humans. *Biosci Biotechnol Biochem* 2001;65:2033 - 6.
15. Ohnuki K, Haramizu S, Watanabe T, Yazawa S, Fushiki T. CH-19 sweet, nonpungent cultivar of red pepper, increased body temperature in mice with vanilloid receptors stimulation by capsiate. *J Nutr Sci Vitaminol* 2001;47:295 - 8.
16. a) Ohnuki K, Haramizu S, Oki K, Watanabe T, Yazawa S, Fushiki T. Administration of capsiate, a nonpungent capsaicin analog, promotes energy metabolism and suppresses body fat accumulation in mice. *Biosci Biotechnol Biochem* 2001;65:2735 - 40.
16. (b) Haramizu S, Mizunoya M, Masuda Y, Ohnuki K, Watanabe T, Yazawa S, Fushiki T. capsiate, a nonpungent capsaicin analog, increases endurance swimming capacity of mice by stimulation of vanilloid receptor. *Biosci Biotechnol Biochem* 2006;70:774 - 81.
17. Lim K, Yoshida M, Kikuzata S, Kiyonaga A, Tanaka H, Shindo M. Dietary red pepper ingestion increase carbohydrate oxidation at rest and during exercise in runners. *Med Sci Sports and Exercise* 1997;29:255 – 361.
18. Kobayashi A, Osaka T, Namba Y, Inoue S, Lee TH, Kimura S. Capsaicin activates heat loss and heat production simultaneously and independently in rats. *Am J Physiol* 1998; 275:R92 - R98.
19. Rosa A, Atzeri A, Deiana M, et al. Protective effect of vanilloids against *tert*-butyl hydroperoxide-induced oxidative stress in *vero* cells culture. *J Agric Food Chem* 2008;56: 3546 - 53.
20. Rosa A, Deiana M, Casu V, Paccagnini S, Appendino G, Dessi MA. Antioxidant activity of capsinoids. *J Agric Food Chem* 2002;50:7396 - 401.
21. Sancho R, Lucena C, Calzado MA, Blanco-Molina M, Minassi A, Appendino G, Munoz E. Immunosuppressive activity of capsaicinoids: capsiate derived from sweet peppers inhibits NF- $\kappa$ B activation and is a potent antiinflammatory compound *in vivo*. *Eur J of Immunology* 2002;32:1753 - 63.

22. Macho A, Lucena C, Sancho R, Daddario N, Minassi A, Munoz E, Appendino G. Non-pungent capsaicinoids from sweet peppers: synthesis and evaluation of the chemopreventive and anticancer potential. *Eur J Nutr* 2003;42:2 - 9.
23. Singh S, Jarret R, Russo V, Majetich G, Shimkus J, Bushway R., Perkins B. Determination of capsinoids by HPLC-DAD in *Capsicum* species. *J Agric Food Chem* 2009;57:3452 - 7.
24. Sutoh K, Kobata K, Yazawa S, Watanabe T. Capsinoids is biosynthesized from phenylalanine and valine in a nonpungent pepper, *Capsicum annuum* L. cv. CH-19 sweet. *Biosci Biotechnol Biochem* 2006;70:1513 - 6.
25. Lang Y, Kisaka H, Sugiyama R, et al. Functional loss of pAMT results in biosynthesis of capsinoids, capsaicinoids analogs, in *Capsicum annuum* cv. CH-19 sweet. *The Plant Journal* 2009;59:953 - 61.
26. Lang Y, Yanagawa S, Sasanuma T, Sasakuma T. Acyl-transferase gene-a candidate gene involved in pungency in capsicum. *Breeding Sci* 2006;54:33 - 9.
27. Stewart C, Kang BC, Liu K, et al. The pun1 gene for pungency in pepper encodes a putative acyltransferase. *Plant J* 2005;42:675 - 88.
28. Stewart C, Mazourek M, Stellari GM, O'Connell M, Jahn MM. Genetic control of pungency in *C. Chinense* via the pun 1 locus. *J Exp Bot* 2007;58:979 - 91.
29. Tanaka Y, Hosokawa M, Miwa T, Watanabe T, Yazawa S. Newly mutated putative-aminotransferase in nonpungent pepper (*Capsicum annuum*) results in biosynthesis of capsinoids, capsaicinoids analogues. *J Agric Food Chem* 2010;58:1761 - 7.
30. Kaga H, Goto K, Takahashi T, Hino M. A general and stereoselective of the capsaicinoids via the orthoester claisen rearrangement. *Tetrahedron* 1996;52:8451 - 70.
31. Kobata K, Kawaguchi M, Watanabe T. Enzymatic synthesis of a capsinoids by the acylation of vanillyl alcohol with fatty acid derivatives catalyzed by lipases. *Biosci Biotechnol Biochem* 2002;66:319 - 27.
32. Ishihara K, Kwon SI, Masuoka N. One-procedure synthesis of capsiate from capsaicin by lipase-catalyzed dynamic transacylation. *World J Microbiol Biotechnol* 2010;26:1337 - 40.
33. [www.centerformedicalweightloss.com/blog/tag/capsiate-natura](http://www.centerformedicalweightloss.com/blog/tag/capsiate-natura) 10/18/2010
34. <http://calorielab.com/news/2010/10/10/capsiate-natura/> 10/18/2010
35. Application for the Approval of Dihydrocapsiate for Use as a Novel Food Ingredient, Ajinomoto CO., INC., JAPAN July 2010.