

## เอกสารอ้างอิง\_หญ้าหนวดแมว

1. ก่องกานดา ชยามฤต. สมุนไพรไทย ตอนที่ 6. กรุงเทพฯ: หจก. ไดมอนด์ พรินติ้ง จำกัด, 2540.
2. Saidan NH, Aisha AF, Hamil MS, et al. A novel reverse phase high-performance liquid chromatography method for standardization of *Orthosiphon stamineus* leaf extracts. *Pharmacog Res.* 2015;7:23-31.
3. Ameer OZ, Salman IM, Asmawi MZ, et al. *Orthosiphon stamineus*: traditional uses, phytochemistry, pharmacology, and toxicology. *J Med Food.* 2012;15:678-90.
4. Olah NK, Radu L, Mogoşan C, et al. Phytochemical and pharmacological studies on *Orthosiphon stamineus* Benth. (Lamiaceae) hydroalcoholic extracts. *J Pharm Biomed Anal.* 2003;33:117-23.
5. Loon YH, Wong JW, Yap SP, et al. Determination of flavonoids from *Orthosiphon stamineus* in plasma using a simple HPLC method with ultraviolet detection. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2005;816:161-6.
6. Ahamed MB, Aisha AF, Nassar ZD, et al. Cat's whiskers tea (*Orthosiphon stamineus*) extract inhibits growth of colon tumor in nude mice and angiogenesis in endothelial cells via suppressing VEGFR phosphorylation. *Nutr Cancer.* 2012;64:89-99.
7. Hussain K, Khan MT, Ismail Z, et al. Rapid separation and determination of betulinic acid from a complex matrix using combination of TLC and RP-HPLC. *Pak J Pharm Sci.* 2012; 25:413-22.
8. Yam MF, Mohamed EA, Ang LF, et al. A simple isocratic HPLC method for the simultaneous determination of sinensetin, eupatorin, and 3'-hydroxy-5,6,7,4'-tetramethoxyflavone in *Orthosiphon stamineus* extracts. *J Acupunct Meridian Stud.* 2012;5:176-82.
9. Mohamed EAH, Mohamed AJ, Asmawi MZ, Sadikun A, Ebrika OS, Yam MF. Antihyperglycemic effect of *Orthosiphon stamineus* Benth leaves extract and its bioassay-guided fractions. *Molecules.* 2011;16:3787-801.
10. Akowuah GA1, Zhari I. Determination of betulinic acid from *Orthosiphon stamineus* leaf extract in rat plasma. *Pharmazie.* 2008;63:788-90.
11. Trimarco V, Cimmino CS, Santoro M, Pagnano G, Manzi MV, Piglia A, et al. Nutraceuticals for blood pressure control in patients with high-normal or grade 1 hypertension. *High Blood Press Cardiovasc Prev.* 2012;19(3):117-22.
12. Werkström V, Andersson KE. ATP- and adenosine-induced relaxation of the smooth muscle of the pig urethra. *BJU Int.* 2005;96:1386-91.
13. Yuliana ND, Khatib A, Link-Struensee AM, et al. Adenosine A1 receptor binding activity of methoxyflavonoids from *Orthosiphon stamineus*. *Planta Med.* 2009;75:132-6.
- 14.

15. Prakasam HS, Herrington H, Roppolo JR, et al. Modulation of bladder function by luminal adenosine turnover and A1 receptor activation. *Am J Physiol Renal Physiol.* 2012;303:F279–92.
16. Adama Y, Somchit MN, Sulaimana,MR, Nasaruddinc AA, Zuraini Ab, Bustamam AA. Diuretic properties of *Orthosiphon stamineus* Benth. *J Ethnopharmacol.* 2009;124:154-8.
17. Beaux D, Fleurentin J, Mortier F. Effect of extracts of *Orthosiphon stamineus* Benth, *Hieracium pilosella* L., *Sambucus nigra* L. and *Arctostaphylos uva-ursi* (L.) Spreng. In rats. *Phytother Res.* 1999;13:222-5.
18. Arafat OM, Thama SY, Sadikuna A, Zharia I, Haughtonb PJ, Asmawia MZ. Studies on diuretic and hypouricemic effects of *Orthosiphon stamineus* methanol extracts in rats. *J Ethnopharmacol.* 2008;118:354-60.
19. Zhong Y-S, Yu C-H, Ying H-Z, Wanga Z-Y, Cai H-F. Prophylactic effects of *Orthosiphon stamineus* Benth. extracts on experimental induction of calcium oxalate nephrolithiasis in rats. *J Ethnopharmacol.* 2012;144:761-7.
20. Laavola M, Nieminen R, Yam MF, Sadikun A, Asmawi MZ, Basir R, et al. Flavonoids eupatorin and sinensetin present in *Orthosiphon stamineus* leaves inhibit inflammatory gene expression and STAT1 activation. *Planta Med.* 2012;78:779-86.
21. Arnold E, Benz T, Zapp C, Wink M. Inhibition of cytosolic phospholipase A<sub>2</sub> $\alpha$  (cPLA<sub>2</sub> $\alpha$ ) by medicinal plants in relation to their phenolic content. *Molecules.* 2015;20:15033-48.
22. Mohamed EAH, Siddiqui MJA, Ang LF, Sadikun A, Chan SH, Tan SC. Potent  $\alpha$ -glucosidase and  $\alpha$ -amylase inhibitory activities of standardized 50% ethanolic extracts and sinensetin from *Orthosiphon stamineus* Benth. as anti-diabetic mechanism. *BMC Complement Altern Med.* 2012;12:176.
23. Mohamed EAH, Yam MF, Ang LF, Mohamed AJ, Asmawi MZ. Antidiabetic properties and mechanism of action of *Orthosiphon stamineus* Benth. bioactive sub-fraction in streptozotocin-induced diabetic rats. *J Acupunct Meridian Stud.* 2013;6(1):31-40.
24. Ngo YL, Chua LS. Anti-diabetic activity of rosmarinic acid rich fractions from *Orthosiphon stamineus*. *Current Enzyme Inhibition.* 2018;14(2):97-103.
25. Guang L; Xi C; Juan L; Xue-lan L; Yi-hang L. Hypoglycemic activity of *Orthosiphon stamineus* aqueous extract on diabetic rats induced by streptozotocin and its mechanism. *Zhonghua Zhongyiyao Zazhi.* 2013;28(12):3653-6.
26. Azam AA, Pariyani R, Ismail IS, Ismail A, Khatib A, Abas F and Shaari K. Urinary metabolomics study on the protective role of *Orthosiphon stamineus* in streptozotocin induced diabetes mellitus in rats via <sup>1</sup>H NMR spectroscopy. *BMC Complement Altern Med.* 2017;17:278
27. Lokman EF, Saparuddina F, Muhammad H, Omard MH, Zulkaplie A. *Orthosiphon stamineus* as a potential antidiabetic drug in maternal hyperglycemia in streptozotocin-induced diabetic rats. *Integr Med Res.* 2019;8:173-9.

28. Yuniarto A, Purwani H, Juanda D, Setiawan F, Kurnia I. Kumis kucing (*Orthosiphon stamineus* Benth.) leaves ethanol extract as anti-obesity agent in high-fat diet-induced obese mice. *Asian J Pharm Clin Res.* 2015;8(6):234-6.
29. Yam MF, Ang LF, Basir R, Salman IM, Ameer OZ, Asmawi MZ. Evaluation of the anti-pyretic potential of *Orthosiphon stamineus* Benth. standardized extract. *Inflammopharmacol.* 2009;17:50-4.
30. Al-Suede FSR, Ahamed MBK, Majid ASA, et al. Optimization of cat's whiskers tea (*Orthosiphon stamineus*) using supercritical carbon dioxide and selective chemotherapeutic potential against prostate cancer cells. *Evid Based Complement Altern Med.* 2014;2014:396016.
31. Dolečková I, Rárová L, Grúz J, et al. Antiproliferative and antiangiogenic effects of flavone eupatorin, an active constituent of chloroform extract of *Orthosiphon stamineus* leaves. *Fitoterapia.* 2012;83:1000-7.
32. Pariyani R, Ismail IS, Azam AA, Abas F, Shaari K, Sulaiman MR. Phytochemical screening and acute oral toxicity study of Java tea leaf extracts. *BioMed Res Int.* 2015:1-8. (Article ID 742420)
33. Purwantiningsih, Hussin AH. Interaction study: the effect of *Orthosiphon stamineus* extract on human cytochrome p450. *Indonesian J Pharm.* 2014;25(4):230-6.
34. กล้วยน้ำว้า : 2.5 ยารักษาอาการทางระบบทางเดินปัสสาวะ. Available from: <http://www.ratchakitcha.soc.go.th/DATA/PDF/2554/E/072/57.PDF> บัญชียาหลัก. [cited 2019 October 28].