

เอกสารอ้างอิง_หอมใหญ่ 38(2)

1. Wound healing and wound care. [อินเทอร์เน็ต]. [สืบค้นเมื่อวันที่ 1 พ.ค. 2563]; จาก: <file:///D:/%E0%B8%AB%E0%B8%B1%E0%B8%A7%E0%B8%AB%E0%B8%AD%E0%B8%A1/use/1.Wound.Healing.pdf>
2. กระบวนการหายของแผลและหลักการรักษา. [อินเทอร์เน็ต]. [สืบค้นเมื่อวันที่ 1 พ.ค. 2563]; จาก: http://202.28.95.4/library/main/eproceeding/Lec_10_17.pdf
3. การดูแลรักษาแผลเป็น (Scar Management). [อินเทอร์เน็ต]. [สืบค้นเมื่อวันที่ 1 พ.ค. 2563]; จาก: <https://he02.tci-thaijo.org/index.php/rtamedj/article/view/61099/50345>
4. แผลเป็น. [อินเทอร์เน็ต]. [สืบค้นเมื่อวันที่ 1 พ.ค. 2563]; จาก: <https://www.si.mahidol.ac.th/sidoctor/e-pl/articledetail.asp?id=548>
5. Bylka W, Znajdek-Awizen P, Studzinska-Sroka E, Brzezinska M. *Centella asiatica* in cosmetology. *Postepy Dermatol Alergol.* 2013;30(1):46-9.
6. Mehta M, Branford OA, Rolfe KJ. The evidence for natural therapeutics as potential anti-scarring agents in burn-related scarring. *Burns Trauma.* 2016;4:15.
7. Song JY, Truong DV, Yang BS. Quercetin shows the pharmacological activity to simultaneously downregulate the inflammatory and fibrotic responses to tissue injury in association with its ability to target multi-kinases. *Pharmacology.* 2018;102(3-4):142-53.
8. Cho JW, Cho SY, Lee SR, Lee KS. Onion extract and quercetin induce matrix metalloproteinase-1 *in vitro* and *in vivo*. *Int J Mol Med.* 2010;25(3):347-52.
9. Phan TT, See P, Tran E, Nguyen TT, Chan SY, Lee ST, et al. Suppression of insulin-like growth factor signalling pathway and collagen expression in keloid-derived fibroblasts by quercetin: its therapeutic potential use in the treatment and/or prevention of keloids. *Br J Dermatol.* 2003;148(3):544-52.
10. Phan TT, Lim IJ, Chan SY, Tan EK, Lee ST, Longaker MT. Suppression of transforming growth factor beta/smad signaling in keloid-derived fibroblasts by quercetin: implications for the treatment of excessive scars. *J Trauma.* 2004;57(5):1032-7.
11. Yuan Z, Yao F, Hu Z, Sun S, Wu B. Quercetin inhibits the migration and proliferation of astrocytes in wound healing. *Neuroreport.* 2015;26(7):387-93.
12. Prager W, Gauglitz GG. Effectiveness and safety of an overnight patch containing *Allium cepa* extract and allantoin for post-dermatologic surgery scars. *Aesthetic Plast Surg.* 2018;42(4):1144-50.
13. Campanati A, Savelli A, Sandroni L, Marconi B, Giuliano A, Giuliodori K, et al. Effect of *Allium cepa*-allantoin-pentaglycan gel on skin hypertrophic scars: clinical and video-capillaroscopic results of an open-label, controlled, nonrandomized clinical trial. *Dermatol Surg.* 2010;36(9):1439-44.

14. Draelos ZD. The ability of onion extract gel to improve the cosmetic appearance of postsurgical scars. *J Cosmet Dermatol*. 2008;7(2):101-4.
15. Draelos ZD, Baumann L, Fleischer AJ, Plaum S, Avakian EV, Hardas B. A new proprietary onion extract gel improves the appearance of new scars: a randomized, controlled, blinded-investigator study. *J Clin Aesthet Dermatol*. 2012;5(6):18-24.
16. Jenwitheesuk K, Surakunprapha P, Jenwitheesuk K, Kuptarnond C, Prathanee S, Intanoo W. Role of silicone derivative plus onion extract gel in presternal hypertrophic scar protection: a prospective randomized, double blinded, controlled trial. *Int Wound J*. 2012;9(4):397-402.
17. Wananukul S, Chatpreodprai S, Peongsujarit D, Lertsapcharoen P. A prospective placebo-controlled study on the efficacy of onion extract in silicone derivative gel for the prevention of hypertrophic scar and keloid in median sternotomy wound in pediatric patients. *J Med Assoc Thai*. 2013;96(11):1428-33.
18. Song T, Kim KH, Lee KW. Randomised comparison of silicone gel and onion extract gel for post-surgical scars. *J Obstet Gynaecol*. 2018;38(5):702-7.
19. Ho WS, Ying SY, Chan PC, Chan HH. Use of onion extract, heparin, allantoin gel in prevention of scarring in Chinese patients having laser removal of tattoos: a prospective randomized controlled trial. *Dermatol Surg*. 2006;32(7):891-6.
20. Koc E, Arca E, Surucu B, Kurumlu Z. An open, randomized, controlled, comparative study of the combined effect of intralesional triamcinolone acetonide and onion extract gel and intralesional triamcinolone acetonide alone in the treatment of hypertrophic scars and keloids. *Dermatol Surg*. 2008;34(11):1507-14.
21. Chanprapaph K, Tanrattanakorn S, Wattanakrai P, Wongkitisophon P, Vachiramon V. Effectiveness of onion extract gel on surgical scars in Asians. *Dermatol Res Pract*. 2012;2012:212945. doi:10.1155/2012/212945.
22. Jackson BA, Shelton AJ. Pilot study evaluating topical onion extract as treatment for postsurgical scars. *Dermatol Surg*. 1999;25(4):267-9.
23. Chung VQ, Kelley L, Marra D, Jiang SB. Onion extract gel versus petrolatum emollient on new surgical scars: prospective double-blinded study. *Dermatol Surg*. 2006;32(2):193-7.