

เอกสารอ้างอิง : กระเทียมบ่มสกัด

1. กองโรคไม่ติดต่อ กรมควบคุมโรค กระทรวงสาธารณสุข. รายงานสถานการณ์โรค NCDs เบาหวาน ความดันโลหิตสูง และปัจจัยที่เกี่ยวข้อง พ.ศ. 2562. [อินเทอร์เน็ต]. 2559. [เข้าถึงเมื่อ 1 กันยายน 2565]. เข้าถึงจาก: <http://www.thaincd.com/document/file/download/knowledge/รายงานสถานการณ์โรคNCDs63update.pdf>
2. Lawson LD. Bioactive organosulfur compounds of garlic and garlic products. Human Medicinal Agents from Plants Chapter 21. American Chemical Society: ACS Symposium Series. doi: 10.1021/bk-1993-0534.ch021.
3. Kang OJ. Physicochemical Characteristics of Black Garlic after Different Thermal Processing Steps. Prev Nutr Food Sci. 2016;21(4):348-354. doi: 10.3746/pnf.2016.21.4.348.
4. Vlachojannis C, Schoenenberger AW, Erne P, Chrubasik-Hausmann S. Preliminary evidence of the clinical effectiveness of odourless garlic. Phytother Res. 2019;33(9):2179-2191. doi: 10.1002/ptr.6409.
5. Kodera Y, Kurita M, Nakamoto M, Matsutomo T. Chemistry of aged garlic: Diversity of constituents in aged garlic extract and their production mechanisms via the combination of chemical and enzymatic reactions. Exp Ther Med. 2020;19(2):1574-84. doi: 10.3892/etm.2019.8393.
6. Abe K, Hori Y, Myoda T. Characterization of key aroma compounds in aged garlic extract. Food Chem. 2020;312:126081. doi: 10.1016/j.foodchem.2019.126081.
7. Kodera Y, Suzuki A, Imada O, Kasuga S, Sumioka I, Kanezawa A, et al. Physical, chemical, and biological properties of S-allylcysteine, an amino acid derived from garlic. J Agric Food Chem. 2002;50(3):622-32. doi: 10.1021/jf0106648.
8. Imai J, Ide N, Nagae S, Moriguchi T, Matsuura H, Itakura Y. Antioxidant and radical scavenging effects of aged garlic extract and its constituents. Planta Med. 1994;60(5):417-20. doi: 10.1055/s-2006-959522.
9. Borek C. Antioxidant health effects of aged garlic extract. J Nutr. 2001;131(3s):1010S-5S. doi: 10.1093/jn/131.3.1010S.
10. Colín-González AL, Santana RA, Silva-Islas CA, Chánez-Cárdenas ME, Santamaría A, Maldonado PD. The antioxidant mechanisms underlying the aged garlic extract- and S-allylcysteine-induced protection. Oxid Med Cell Longev. 2012;2012:907162. doi: 10.1155/2012/907162.
11. Nagae S, Ushijima M, Hatono S, Imai J, Kasuga S, Matsuura H, Itakura Y, et al. Pharmacokinetics of the garlic compound S-allylcysteine. Planta Med. 1994;60(3):214-7. doi: 10.1055/s-2006-959461.

12. Budoff MJ, Takasu J, Flores FR, Niihara Y, Lu B, Lau BH, et al. Inhibiting progression of coronary calcification using Aged Garlic Extract in patients receiving statin therapy: a preliminary study. *Prev Med.* 2004;39(5):985-91. doi: 10.1016/j.ypmed.2004.04.012.
13. Wlosinska M, Nilsson AC, Hlebowicz J, Hauggaard A, Kjellin M, Fakhro M, et al. The effect of aged garlic extract on the atherosclerotic process - a randomized double-blind placebo-controlled trial. *BMC Complement Med Ther.* 2020;20(1):132. doi: 10.1186/s12906-020-02932-5.
14. Matsumoto S, Nakanishi R, Li D, Alani A, Rezaeian P, Prabhu S, et al. Aged garlic extract reduces low attenuation plaque in coronary arteries of patients with metabolic syndrome in a prospective randomized double-blind study. *J Nutr.* 2016;146(2):427S-32S. doi: 10.3945/jn.114.202424.
15. Budoff MJ, Ahmadi N, Gul KM, Liu ST, Flores FR, Tiano J, et al. Aged garlic extract supplemented with B vitamins, folic acid and L-arginine retards the progression of subclinical atherosclerosis: a randomized clinical trial. *Prev Med.* 2009;49(2-3):101-7. doi: 10.1016/j.ypmed.2009.06.018.
16. Ahmadi N, Tsimikas S, Hajsadeghi F, Saeed A, Nabavi V, Bevin MA, et al. Relation of oxidative biomarkers, vascular dysfunction, and progression of coronary artery calcium. *Am J Cardiol.* 2010;105(4):459-66. doi: 10.1016/j.amjcard.2009.09.052.
17. Ahmadi N, Nabavi V, Hajsadeghi F, Zeb I, Flores F, Ebrahimi R, et al. Aged garlic extract with supplement is associated with increase in brown adipose, decrease in white adipose tissue and predict lack of progression in coronary atherosclerosis. *Int J Cardiol.* 2013; 168(3):2310-4. doi: 10.1016/j.ijcard.2013.01.182.
18. Zeb I, Ahmadi N, Flores F, Budoff MJ. Randomized trial evaluating the effect of aged garlic extract with supplements versus placebo on adipose tissue surrogates for coronary atherosclerosis progression. *Coron Artery Dis.* 201;29(4):325-8. doi: 10.1097/MCA.0000000000000587.
19. Weiss N, Ide N, Abahji T, Nill L, Keller C, Hoffmann U. Aged garlic extract improves homocysteine-induced endothelial dysfunction in macro- and microcirculation. *J Nutr.* 2006;136(3 Suppl):750S-4S.
20. Gruenwald J, Bongartz U, Bothe G, Uebelhack R. Effects of aged garlic extract on arterial elasticity in a placebo-controlled clinical trial using EndoPATTM technology. *Exp Ther Med.* 2020;19(2):1490-1499. doi: 10.3892/etm.2019.8378.
21. Williams MJ, Sutherland WH, McCormick MP, Yeoman DJ, de Jong SA. Aged garlic extract improves endothelial function in men with coronary artery disease. *Phytother Res.* 2005; 19(4):314-9. doi: 10.1002/ptr.1663.

22. Larijani VN, Ahmadi N, Zeb I, Khan F, Flores F, Budoff M. Beneficial effects of aged garlic extract and coenzyme Q10 on vascular elasticity and endothelial function: the FAITH randomized clinical trial. *Nutrition*. 2013;29(1):71-5. doi: 10.1016/j.nut.2012.03.016.
23. Rahman K, Billington D. Dietary supplementation with aged garlic extract inhibits ADP-induced platelet aggregation in humans. *J Nutr*. 2000;130(11):2662-5. doi: 10.1093/jn/130.11.2662.
24. Steiner M, Li W. Aged garlic extract, a modulator of cardiovascular risk factors: a dose-finding study on the effects of AGE on platelet functions. *J Nutr*. 2001;131(3s):980S-4S. doi: 10.1093/jn/131.3.980S.
25. Lindstedt S, Wlosinska M, Nilsson AC, Hlebowicz J, Fakhro M, Sheikh R. Successful improved peripheral tissue perfusion was seen in patients with atherosclerosis after 12 months of treatment with aged garlic extract. *Int Wound J*. 2021;18(5):681-691. doi: 10.1111/iwj.13570.
26. Wlosinska M, Nilsson AC, Hlebowicz J, Malmsjö M, Fakhro M, Lindstedt S. Aged garlic extract preserves cutaneous microcirculation in patients with increased risk for cardiovascular diseases: A double-blinded placebo-controlled study. *Int Wound J*. 2019;16(6):1487-93. doi: 10.1111/iwj.13220.
27. Ried K, Frank OR, Stocks NP. Aged garlic extract reduces blood pressure in hypertensives: a dose-response trial. *Eur J Clin Nutr*. 2013;67(1):64-70. doi: 10.1038/ejcn.2012.178.
28. Ried K, Frank OR, Stocks NP. Aged garlic extract lowers blood pressure in patients with treated but uncontrolled hypertension: a randomised controlled trial. *Maturitas*. 2010;67(2):144-50. doi: 10.1016/j.maturitas.2010.06.001.
29. Steiner M, Khan AH, Holbert D, Lin RI. A double-blind crossover study in moderately hypercholesterolemic men that compared the effect of aged garlic extract and placebo administration on blood lipids. *Am J Clin Nutr*. 1996;64(6):866-70. doi: 10.1093/ajcn/64.6.866.
30. Seo DY, Lee SR, Kim HK, Baek YH, Kwak YS, Ko TH, et al. Independent beneficial effects of aged garlic extract intake with regular exercise on cardiovascular risk in post-menopausal women. *Nutr Res Pract*. 2012;6(3):226-31. doi: 10.4162/nrp.2012.6.3.226.
31. Amagase H, Petesch BL, Matsuura H, Kasuga S, Itakura Y. Intake of garlic and its bioactive components. *J Nutr*. 2001;131(3s):955S-62S. doi: 10.1093/jn/131.3.955S.
32. Macan H, Uykimpang R, Alconcel M, Takasu J, Razon R, Amagase H, et al. Aged garlic extract may be safe for patients on warfarin therapy. *J Nutr*. 2006;136:793S-5S.
33. กฤติยา ไชยนอก. อันตรกิริยาของกระเทียมกับยาแผนปัจจุบัน. *จุลสารข้อมูลสมุนไพร*. 2562;37(1):12-24.