

เอกสารอ้างอิง : เทียนเกล็ดหอย 41(3)

1. นันทวัน บุญยะประภัศร และอรนุช โชคชัยเจริญพร, บรรณาธิการ. สมุนไพร..ไม้พื้นบ้าน (2). กรุงเทพฯ: บริษัท ประชาชน จำกัด; 2541.
2. ชัยนัต พิเชียรสุนทร แม้นมาส ชวลิต วิเชียร จีรวงศ์. คำอธิบายตำราพระโอสถพระนารายณ์. กรุงเทพมหานคร: สำนักพิมพ์อมรินทร์; 2542.
3. *Plantago ovata* Forssk. World Flora Online. [Internet]. 2012 [cited 2022 Dec 7]. Available from: <http://www.worldfloraonline.org/taxon/wfo-0000486655>
4. นพมาศ สุนทรเจริญนนท์ และนางลักษณ์ เรืองวิเศษ. คุณภาพเครื่องยาไทยจากงานวิจัยสู่การพัฒนาอย่างยั่งยืน. กรุงเทพฯ: สำนักงานคณะกรรมการวิจัยแห่งชาติ; 2551.
5. กองผลิตภัณฑ์สมุนไพร. ประกาศคณะกรรมการพัฒนาระบบยาแห่งชาติ เรื่อง บัญชียาหลักแห่งชาติ ด้านสมุนไพร พ.ศ. 2566 [อินเทอร์เน็ต]. นนทบุรี: สำนักงานคณะกรรมการอาหารและยา; 2566 [เข้าถึงเมื่อ 18 ก.ค. 2566]. เข้าถึงได้จาก: <https://ratchakitcha.soc.go.th/documents/140D130S0000000004500.pdf>
6. Der Marderosian A, editor. The Review of Natural Products. The most complete source of natural product information 2001. 1st ed. Missouri: Facts and Comparisons; 2001.
7. Edwards CA, Bowen J, Brydon WG, Eastwood MA. The effects of ispaghula on rat caecal fermentation and stool output. Br J Nutr. 1992;68(2):473-82.
8. Leng-Peschlow E. *Plantago ovata* seeds as dietary fibre supplement: physiological and metabolic effects in rats. Br J Nutr. 1991;66(2):331-49.
9. Yokota M, Mochizuki Y, Mori Y, Kanbe H. Constipation therapeutic agents comprising *Plantago ovata* testae. Repub. Korean Kongkae Taeho Kongbo 2010 patent KR 2010036954 A 20100408.
10. Rodríguez-Cabezas ME, Gáálvez J, Lorente MD, Concha A, Camuesco D, Azzouz S, et al. Dietary fiber down-regulates colonic tumor necrosis factor alpha and nitric oxide production in trinitrobenzenesulfonic acid-induced colitic rats. J Nutr. 2002;132(11):3263-71.
11. Rodríguez-Cabezas ME, Gáálvez J, Camuesco D, Lorente MD, Concha A, Martínez-Augustín O, Redondo L, Zarzuelo A. Intestinal anti-inflammatory activity of dietary fiber (*Plantago ovata* seeds) in HLA-B27 transgenic rats. Clin Nutr. 2003;22(5):463-71.
12. Ewe K, Ueberschaer B, Press AG. Influence of senna, fibre, and fibre + senna on colonic transit in loperamide-induced constipation. Pharmacology. 1993;47Suppl 1:242-8.
13. Tomás-Ridocci M, Añóñón R, Miñiguez M, Zaragoza A, Ballester J, Benages A. The efficacy of *Plantago ovata* as a regulator of intestinal transit. A double-blind study compared to placebo. Rev Esp Enferm Dig. 1992;82(1):17-22.
14. Marlett JA, Kajs TM, Fischer MH. An unfermented gel component of psyllium seed husk promotes laxation as a lubricant in humans. Am J Clin Nutr. 2000;72(3):784-89.

15. Galisteo M, Sáánchez M, Vera R, Gonzáález M, Anguera A, Duarte J, Zarzuelo A. A diet supplemented with husks of *Plantago ovata* reduces the development of endothelial dysfunction, hypertension, and obesity by affecting adiponectin and TNF-alpha in obese Zucker rats. *J Nutr.* 2005;135(10):2399-404.
16. Miettinen TA, Tarpila S. Serum lipids and cholesterol metabolism during guar gum, *Plantago ovata* and high fibre treatments. *Clin Chim Acta.* 1989;183(3):253-62.
17. Romero AL, West KL, Zern T, Fernandez ML. The seeds from *Plantago ovata* lower plasma lipids by altering hepatic and bile acid metabolism in guinea pigs. *J Nutr.* 2002;132(6):1194-8.
18. Ulbricht CE, Basch EM, editors. *Natural Standard Herb & Supplement Reference: Evidence-based Clinical Reviews.* Missouri: Elsevier Mosby; 2005.
19. Moreno LA, Tresaco B, Bueno G, Fleta J, Rodríguez G, Garagorri JM, et al. Psyllium fibre and the metabolic control of obese children and adolescents. *J Physiol Biochem.* 2003;59(3):235-42.
20. Sola R, Goda G, Ribalta J, Vallvé J-C, Girona J, Anguera A, et al. Effects of soluble fiber (*Plantago ovata* husk) on plasma lipids, lipoproteins, and apolipoproteins in men with ischemic heart disease. *Am J Clin Nutr.* 2007;85(4):1157-63.
21. Uehleke B, Ortiz M, Stange R. Cholesterol reduction using psyllium husks – Do gastrointestinal adverse effects limit compliance? Results of a specific observational study. *Phytomedicine.* 2008;15:153-9.
22. Pal S, Khossousi A, Binns C, Dhaliwal S, Ellis V. The effect of a fibre supplement compared to a healthy diet on body composition, lipids, glucose, insulin and other metabolic syndrome risk factors in overweight and obese individuals. *Br J Nutr.* 2011; 105:90-100.
23. Khossousi A, Binns CW, Dhaliwal SS, Pal S. The acute effects of psyllium on postprandial lipaemia and thermogenesis in overweight and obese men. *Br J Nutr.* 2008;99:1068-75.
24. Salas-Salvadó J, Farrés X, Luque X, Narejos S, Borrell M, Basora J, et al. Effect of two doses of a mixture of soluble fibres on body weight and metabolic variables in overweight or obese patients: A randomised trial. *Br J Nutr.* 2008;99(6):1380-7.
25. Howard AN, Marks J. Hypocholesterolemic effect of dietary fibers: studies on the mechanism of action. *Arterioscler Brain Dis.* 1983:203-5.
26. Hannan JM, Ali L, Khaleque J, Akhter M, Flatt PR, Abdel-Wahab YH. Aqueous extracts of husks of *Plantago ovata* reduce hyperglycaemia in type 1 and type 2 diabetes by inhibition of intestinal glucose absorption. *Br J Nutr.* 2006;96(1):131-7.
27. Ziai SA, Larijani B, Akhoondzadeh S, Fakhrzadeh H, Dastpak A, Bandarian F, Rezai A, Badi HN, Emami T. Psyllium decreased serum glucose and glycosylated hemoglobin significantly in diabetic outpatients. *J Ethnopharmacol.* 2005;102(2):202-7.

28. Machado L, Zetterström O, Fagerberg E. Occupational allergy in nurses to a bulk laxative. *Allergy*. 1979;34(1):51-5.
29. Suhonen R, Kantola I, Björkstén F. Anaphylactic shock due to ingestion of psyllium laxative. *Allergy*. 1983;38(5):363-5.
30. Freeman GL. Psyllium hypersensitivity. *Ann Allergy* 1994; 73(6): 490-2.
31. Shin S, Kim JH, Mun YH, Chung HS. Acute esophageal obstruction after ingestion of psyllium seed husk powder: A case report. *World J Clin Cases*. 2022;10(7):2336-2340. doi: 10.12998/wjcc.v10.i7.2336.
32. Salguero Molpeceres O, Seijas Ruiz-Coello MC, Hernández Núñez J, Caballos Villar D, Díaz Picazo L, Ayerbe García-Monzón L. Esophageal obstruction caused by dietary fiber from *Plantago ovata*, a complication preventable by adequate information. *Gastroenterol Hepatol*. 2003;26(4):248-50.
33. Food and Drug Administration, HHS. Laxative drug products for over-the-counter human use; psyllium ingredients in granular dosage forms. Final rule. *Fed Regis*. 2007; 72(60):14669-74.
34. Robinson DS, Benjamin DM, McCormack JJ. Interaction of warfarin and nonsystemic gastrointestinal drugs. *Clin Pharmacol Ther*. 1971;12(3):491-5. doi: 10.1002/cpt1971123491.
35. Nordström M, Melander A, Robertsson E, Steen B. Influence of wheat bran and of a bulk-forming ispaghula cathartic on the bioavailability of digoxin in geriatric in-patients. *Drug Nutr Interact*. 1987;5(2):67-9.
36. Toutoungi M, Schulz P, Widmer J, Tissot R. Probable interaction entre le psyllium et le lithium [Probable interaction of psyllium and lithium]. *Thérapie*. 1990;45(4):358-60. French.
37. Perlman BB. Interaction between lithium salts and ispaghula husk. *Lancet*. 1990;335(8686):416. doi: 10.1016/0140-6736(90)90256-5.
38. Etman MA. Effect of a bulk forming laxative on the bioavailability of carbamazepine in man. *Drug Dev Ind Pharm*. 1995;21(16):1901-6.
39. Ettinger AB, Shinnar S, Sinnott MJ, Moshe SL. Carbamazepine-induced constipation. *J Epilepsy*. 1992;5(3):191-3. doi: 10.1016/S0896-6974(05)80140-9.
40. Turley SD, Daggy BP, Dietschy JM. Effect of feeding psyllium and cholestyramine in combination on low density lipoprotein metabolism and fecal bile acid excretion in hamsters with dietary-induced hypercholesterolemia. *J Cardiovasc Pharmacol*. 1996; 27(1):71-9. doi: 10.1097/00005344-199601000-00012.
41. Daggy BP, O'Connell NC, Jerdack GR, Stinson BA, Setchell KD. Additive hypocholesterolemic effect of psyllium and cholestyramine in the hamster: influence on fecal sterol and bile acid profiles. *J Lipid Res*. 1997;38(3):491-502.

42. Spence JD, Huff MW, Heidenheim P, Viswanatha A, Munoz C, Lindsay R, et al. Combination therapy with colestipol and psyllium mucilloid in patients with hyperlipidemia. *Ann Intern Med.* 1995;123(7):493-9. doi: 10.7326/0003-4819-123-7-199510010-00003.
43. Cavaliere H, Floriano I, Medeiros-Neto G. Gastrointestinal side effects of orlistat may be prevented by concomitant prescription of natural fibers (psyllium mucilloid). *Int J Obes Relat Metab Disord.* 2001;25(7):1095-9. doi: 10.1038/sj.ijo.0801645.
44. Agrawal AR, Tandon M, Sharma PL. Effect of combining viscous fibre with lovastatin on serum lipids in normal human subjects. *Int J Clin Pract.* 2007;61(11):1812-8. doi: 10.1111/j.1742-1241.2007.01512.x.
45. Garcia JJ, Fernandez N, Carriedo D, Diez MJ, Sahagun A, Gonzalez A, et al. Hydrosoluble fiber (*Plantago ovata* husk) and levodopa I: experimental study of the pharmacokinetic interaction. *Eur Neuropsychopharmacol.* 2005;15(5):497-503. doi: 10.1016/j.euroneuro.2005.01.005.
46. Fernandez N, Carriedo D, Sierra M, Diez MJ, Sahagun A, Calle A, et al. Hydrosoluble fiber (*Plantago ovata* husk) and levodopa II: experimental study of the pharmacokinetic interaction in the presence of carbidopa. *Eur Neuropsychopharmacol.* 2005;15(5):505-9. doi: 10.1016/j.euroneuro.2005.01.006.
47. García JJ, Fernández N, Calle AP, Diez MJ, Sahagún A, Sierra M. Effects of *Plantago ovata* husk on levodopa (with Carbidopa) bioavailability in rabbits with autonomic gastrointestinal disorders. *Drug Metab Dispos.* 2009;37(7):1434-42. doi: 10.1124/dmd.108.026229.
48. Fernandez-Martinez MN, Hernandez-Echevarria L, Sierra-Vega M, Diez-Liebana MJ, Calle-Pardo A, Carriedo-Ule D, et al. A randomised clinical trial to evaluate the effects of *Plantago ovata* husk in Parkinson patients: changes in levodopa pharmacokinetics and biochemical parameters. *BMC Complement Altern Med.* 2014;14:296. doi: 10.1186/1472-6882-14-296.
49. Díez-Láiz R, García-Vieitez JJ, Díez-Liébana MJ, Sierra-Vega M, Sahagún-Prieto AM, Calle-Pardo AP, et al. Evaluation of the association metformin: *Plantago ovata* husk in diabetic rabbits. *J Diabetes Res.* 2015;2015:167526. doi: 10.1155/2015/167526.
50. Díez R, García JJ, Diez MJ, Sierra M, Sahagun AM, Fernández N. Influence of *Plantago ovata* husk (dietary fiber) on the bioavailability and other pharmacokinetic parameters of metformin in diabetic rabbits. *BMC Complement Altern Med.* 2017; 17(1):298. doi: 10.1186/s12906-017-1809-x.
51. Fernández-Bañares F, Hinojosa J, Sánchez-Lombraña JL, Navarro E, Martínez-Salmerón JF, García-Pugés A, et al. Randomized clinical trial of *Plantago ovata* seeds (dietary fiber) as compared with mesalamine in maintaining remission in ulcerative colitis.

Spanish Group for the Study of Crohn's Disease and Ulcerative Colitis (GETECCU). *Am J Gastroenterol.* 1999;94(2):427-33. doi: 10.1111/j.1572-0241.1999.872_a.x.

52. Ahi S, Esmailzadeh M, Kayvanpour E, Sedaghat-Hamedani F, Samadanifard SH. A bulking agent may lead to adrenal insufficiency crisis: a case report. *Acta Med Iran.* 2011;49(10):688-9.
53. Ebadollahi-Natanzi A, Arab-Rahmatipour G. Psyllium together with allopurinol can efficiently reduce the increased serum level of uric acid, creatinine and urea: a case report. *Iran J Toxicol.* 2017;11(4):51-6.